

**12th Romanian-Finnish Seminar, Turku, 19.8.2009**

# **Close-to-convexity of quasihyperbolic balls**

**Riku Klén**  
**riku.klen@utu.fi**

**Abstract.** We will consider close-to-convexity of metric balls defined by the quasihyperbolic metric and the  $j$ -metric.

## Definitions

**Let  $G \subsetneq \mathbb{R}^n$ ,  $n \geq 2$ , be a domain and define**

- **The *quasihyperbolic distance* for  $x, y \in G$  by**

$$k_G(x, y) = \inf_{\alpha \in \Gamma_{xy}} \int_{\alpha} \frac{|dz|}{d(z)},$$

**where  $d(z) = d(z, \partial G)$  and  $\Gamma_{xy}$  is the collection of all rectifiable curves in  $G$  joining  $x$  and  $y$ .**

- **[MO] For  $x, y \in \mathbb{R}^n \setminus \{0\}$  and  $n \geq 2$**

$$k_{\mathbb{R}^n \setminus \{0\}}(x, y) = \sqrt{\alpha^2 + \log^2 \frac{|x|}{|y|}},$$

**where  $\alpha = \angle(x, 0, y) \in [0, \pi]$ .**

- **The  $j$ -distance for  $x, y \in G$  by**

$$j_G(x, y) = \log \left( 1 + \frac{|x - y|}{\min\{d(x), d(y)\}} \right).$$


**Note that  $j_G(x, y) \leq k_G(x, y)$  for all  $x, y \in G$ .**

- **For  $m \in \{k_G, j_G\}$  we define the *metric ball (metric disk* in the case  $n = 2$ ) for  $r > 0$  and  $x \in G$  by**

$$B_m(x, r) = \{y \in G : m(x, y) < r\}.$$

- $G$  is **starlike with respect to**  $x \in G$  if for all  $y \in G$  the line segment  $[x, y]$  is contained in  $G$  and  $G$  is **strictly starlike with respect to**  $x$  if each line from the point  $x$  meets  $\partial G$  at exactly one point.
- If  $G$  is starlike for all  $x \in G$  then it is **convex**.
- A domain  $G \subset \mathbb{R}^n$  is **close-to-convex** if  $\mathbb{R}^n \setminus G$  can be covered with non-intersecting half-lines  
 $(\{z \in \mathbb{R}^n : z = x + ty, x, y \in \mathbb{R}^n, y \neq 0, t > 0\}$  or  
 $\{z \in \mathbb{R}^n : z = x + ty, x, y \in \mathbb{R}^n, y \neq 0, t \geq 0\})$ .

**Clearly convex domains are starlike and starlike domains are close-to-convex.**



**Figure 1: An example of quasihyperbolic and  $j$ -metric disks in a punctured plane.**

## Close-to-convexity

**Let  $D$  be the unit disk and  $f: D \rightarrow D' = f(D)$  be a univalent function with  $f(0) = 0$ . Then**

- **$D'$  is convex  $\iff \operatorname{Re} \frac{zf'(z)}{f(z)} > \frac{1}{2}$  [S, Corollary 2.15].**
- **$D'$  is starlike w.r.t. 0  $\iff \operatorname{Re} \frac{zf'(z)}{f(z)} > 0$  [S, Theorem 2.2].**

**Close-to-convex functions were introduced by W. Kaplan in 1952 [K].**

- **$f(z)$  is close-to-convex iff there exists a convex  $g$  such that  $\operatorname{Re} \frac{f'(z)}{g'(z)} > 0$ .**

**Lewandowski has shown that [S] that  $f$  is close-to-convex iff  $D'$  is close-to-convex.**

## Geometry of the quasihyperbolic balls

The work is based on the following open problem [Vu, 8.1]:

**Does there exist  $r_0 > 0$  such that  $B_m(x, r)$  is convex (in Euclidean geometry) for all  $r \in (0, r_0)$ ? ( $G \subset \mathbb{R}^n$  domain,  $m$  metric)**

**Let  $G \subsetneq \mathbb{R}^n$  be a domain and  $x \in G$ . Then  $B_k(x, r)$  is**

- **convex for  $n = 2$  and  $r \in (0, 1]$  [V],**
- **convex for  $r > 0$ , if  $G$  is convex [MV],**
- **starlike w.r.t  $x$  for  $r > 0$ , if  $G$  is starlike w.r.t.  $x$  [K].**

# References

- [K] R. Klén: ***Local Convexity Properties of Quasihyperbolic Balls in Punctured Space.*** J. Math. Anal. Appl. 342 (2008), 192–201.
- [MO] G.J. Martin, B.G. Osgood: ***The quasihyperbolic metric and the associated estimates on the hyperbolic metric.*** J. Anal. Math. 47 (1986), 37–53.
- [MV] O. Martio and J. Väisälä: ***Quasihyperbolic geodesics in convex domains II.*** To appear in Pure Appl. Math. Q.
- [S] T.J. Suffridge: ***Some special classes of conformal mappings.*** Handbook of complex analysis: geometric function theory. Vol. 2, 309–338, Elsevier, Amsterdam, 2005.
- [V] J. Väisälä: ***Quasihyperbolic geometry of planar domains.*** Manuscript, June, 2008.
- [Vu] M. Vuorinen: **Metrics and quasiregular mappings.** - In Quasiconformal Mappings and their Applications (New Delhi, India, 2007), S. Ponnusamy, T. Sugawa, and M. Vuorinen , Eds., Narosa Publishing House, pp. 291–325.