

# On the characterization of all $\beta$ with $(\pm\beta)$ -integers encoded by conjugated morphisms

Daniel Dombek

*joint work with Z. Masáková and T. Vávra*

Czech Technical University in Prague

WORDS 2013, Turku

17 September 2013

# Table of contents

- 1  $(\pm\beta)$ -integers
- 2 Similarity of  $\mathbb{Z}_\beta^+$  and  $\mathbb{Z}_{-\beta}$
- 3 Cubic Pisot examples

# Table of contents

1  $(\pm\beta)$ -integers

2 Similarity of  $\mathbb{Z}_\beta^+$  and  $\mathbb{Z}_{-\beta}$

3 Cubic Pisot examples

# $b$ -expansions

Let  $b \in \mathbb{R}$ ,  $|b| > 1$ ,  $x \in [\ell, \ell + 1)$ ,  $\ell \in \mathbb{R}$ . The  $(b, \ell)$ -expansion of  $x$  is the sequence  $(x_i)_{i \geq 1}$ , where

$$x_i := \lfloor bT^{i-1}(x) - \ell \rfloor,$$
$$T(x) := bx - \lfloor bx - \ell \rfloor.$$

# $b$ -expansions

Let  $b \in \mathbb{R}$ ,  $|b| > 1$ ,  $x \in [\ell, \ell + 1)$ ,  $\ell \in \mathbb{R}$ . The  $(b, \ell)$ -expansion of  $x$  is the sequence  $(x_i)_{i \geq 1}$ , where

$$\begin{aligned}x_i &:= \lfloor bT^{i-1}(x) - \ell \rfloor, \\T(x) &:= bx - \lfloor bx - \ell \rfloor.\end{aligned}$$

We write  $d(x) = d_{b,\ell}(x) := x_1x_2x_3 \cdots$  and it holds

$$x = \frac{x_1}{b} + \frac{x_2}{b^2} + \frac{x_3}{b^3} + \cdots.$$

## $b$ -expansions $\cdots$ admissibility

$(x_i)_{i \in \mathbb{N}}$  is  $(b, \ell)$ -admissible if  $d(x) = x_1 x_2 \cdots$  for some  $x \in [\ell, \ell + 1)$

# $b$ -expansions $\cdots$ admissibility

$(x_i)_{i \in \mathbb{N}}$  is  $(b, \ell)$ -admissible if  $d(x) = x_1 x_2 \cdots$  for some  $x \in [\ell, \ell + 1)$

## Theorem

For fixed  $b, \ell$  it holds:  $(x_i)_{i \in \mathbb{N}}$  is  $(b, \ell)$ -admissible iff

$$d(\ell) \preceq x_k x_{k+1} x_{k+2} \cdots \prec d^*(\ell + 1) = \lim_{\varepsilon \rightarrow 0^+} d(\ell + 1 - \varepsilon),$$

for all  $k \geq 1$ , where

- $b = \beta > 1 \rightarrow$  lexicographic ordering  $\preceq_{\text{lex}}$
- $b = -\beta < -1 \rightarrow$  alternate ordering  $\preceq_{\text{alt}}$

# $b$ -expansions $\cdots$ admissibility

$(x_i)_{i \in \mathbb{N}}$  is  $(b, \ell)$ -admissible if  $d(x) = x_1 x_2 \cdots$  for some  $x \in [\ell, \ell + 1)$

## Theorem

For fixed  $b, \ell$  it holds:  $(x_i)_{i \in \mathbb{N}}$  is  $(b, \ell)$ -admissible iff

$$d(\ell) \preceq x_k x_{k+1} x_{k+2} \cdots \prec d^*(\ell + 1) = \lim_{\varepsilon \rightarrow 0+} d(\ell + 1 - \varepsilon),$$

for all  $k \geq 1$ , where

- $b = \beta > 1 \rightarrow$  lexicographic ordering  $\preceq_{\text{lex}}$
- $b = -\beta < -1 \rightarrow$  alternate ordering  $\preceq_{\text{alt}}$

$$u_1 u_2 \cdots \prec_{\text{alt}} v_1 v_2 \cdots \text{ if } (-1)^k (u_k - v_k) < 0, \quad k = \min\{i \geq 1, u_i \neq v_i\}$$



# $(\pm\beta)$ -expansions of reals

Rényi  $\beta$ -expansions (1957):

- $b = \beta > 1$ ,  $[\ell, \ell + 1) = [0, 1)$
- $x_i \in \{0, 1, \dots, \lceil \beta \rceil - 1\}$
- Any  $x \in \mathbb{R}$  has a unique  $\beta$ -expansion  
 $(x)_\beta = (\pm)x_k x_{k-1} \cdots x_1 x_0 \bullet x_{-1} x_{-2} \cdots$

# $(\pm\beta)$ -expansions of reals

Rényi  $\beta$ -expansions (1957):

- $b = \beta > 1$ ,  $[\ell, \ell + 1) = [0, 1)$
- $x_i \in \{0, 1, \dots, \lceil \beta \rceil - 1\}$
- Any  $x \in \mathbb{R}$  has a unique  $\beta$ -expansion  
 $(x)_\beta = (\pm)x_k x_{k-1} \cdots x_1 x_0 \bullet x_{-1} x_{-2} \cdots$

Ito-Sadahiro  $(-\beta)$ -expansions (2009):

- $b = -\beta < -1$ ,  $[\ell, \ell + 1) = [\frac{-\beta}{\beta+1}, \frac{1}{\beta+1})$
- $x_i \in \{0, 1, \dots, \lfloor \beta \rfloor\}$
- For any  $x \in \mathbb{R}$  we can define a unique  $(-\beta)$ -expansion  
 $\langle x \rangle_{-\beta} = x_k x_{k-1} \cdots x_1 x_0 \bullet x_{-1} x_{-2} \cdots$

# $(\pm\beta)$ -integers

$\beta$ -integers:

$$\mathbb{Z}_\beta^+ := \{x \in \mathbb{R}^+ : (x)_\beta = x_k \cdots x_1 x_0 \bullet 0^\omega\} = \bigcup_{n \geq 0} \beta^n T_\beta^{-n}(0)$$

$$\mathbb{Z}_\beta := \mathbb{Z}_\beta^+ \cup (-\mathbb{Z}_\beta^+)$$

# $(\pm\beta)$ -integers

$\beta$ -integers:

$$\mathbb{Z}_\beta^+ := \{x \in \mathbb{R}^+ : (x)_\beta = x_k \cdots x_1 x_0 \bullet 0^\omega\} = \bigcup_{n \geq 0} \beta^n T_\beta^{-n}(0)$$

$$\mathbb{Z}_\beta := \mathbb{Z}_\beta^+ \cup (-\mathbb{Z}_\beta^+)$$

$(-\beta)$ -integers:

$$\mathbb{Z}_{-\beta} := \{x \in \mathbb{R} : \langle x \rangle_{-\beta} = x_k \cdots x_1 x_0 \bullet 0^\omega\} = \bigcup_{n \geq 0} (-\beta)^n T_{-\beta}^{-n}(0)$$

## ( $\pm\beta$ )-integers $\cdots$ encoding by infinite word

Let  $\beta > 1$  be a Pisot number (algebraic integer, conjugates  $|\cdot| < 1$ ). Then:

$\mathbb{Z}_\beta^+$  : is an infinite set ,  
contains distances of lengths  $\Delta_0, \dots, \Delta_k$ , all  $\leq 1$  ,  
can be encoded by infinite word  $\mathbf{u}_\beta \in \{0, \dots, k\}^\mathbb{N}$  ,  
there exists a morphism  $\varphi$  fixing  $\mathbf{u}_\beta$  .

## $(\pm\beta)$ -integers $\cdots$ encoding by infinite word

Let  $\beta > 1$  be a Pisot number (algebraic integer, conjugates  $|\cdot| < 1$ ). Then:

$\mathbb{Z}_\beta^+$  : is an infinite set ,

contains distances of lengths  $\Delta_0, \dots, \Delta_k$ , all  $\leq 1$ ,  
can be encoded by infinite word  $\mathbf{u}_\beta \in \{0, \dots, k\}^{\mathbb{N}}$ ,  
there exists a morphism  $\varphi$  fixing  $\mathbf{u}_\beta$ .

$\mathbb{Z}_{-\beta}$  : is an infinite set  $\Leftrightarrow \beta \geq \tau = (1 + \sqrt{5})/2$ ,

contains distances of lengths  $\Delta'_0, \dots, \Delta'_{k'}$ ,  
can be encoded by biinfinite word  $\mathbf{v}_{-\beta} \in \{0, \dots, k'\}^{\mathbb{Z}}$ ,  
there exists an antimorphism  $\psi$  fixing  $\mathbf{v}_{-\beta}$ .

# $(\pm\beta)$ -integers $\cdots$ encoding by infinite word

Let  $\beta > 1$  be a Pisot number (algebraic integer, conjugates  $|\cdot| < 1$ ). Then:

$\mathbb{Z}_\beta^+$  : is an infinite set ,  
contains distances of lengths  $\Delta_0, \dots, \Delta_k$ , all  $\leq 1$  ,  
can be encoded by infinite word  $\mathbf{u}_\beta \in \{0, \dots, k\}^\mathbb{N}$  ,  
there exists a morphism  $\varphi$  fixing  $\mathbf{u}_\beta$  .

$\mathbb{Z}_{-\beta}$  : is an infinite set  $\Leftrightarrow \beta \geq \tau = (1 + \sqrt{5})/2$  ,  
contains distances of lengths  $\Delta'_0, \dots, \Delta'_{k'}$  ,  
can be encoded by biinfinite word  $\mathbf{v}_{-\beta} \in \{0, \dots, k'\}^\mathbb{Z}$  ,  
there exists an antimorphism  $\psi$  fixing  $\mathbf{v}_{-\beta}$  .

- $\varphi(xy) = \varphi(x)\varphi(y)$ ,  $\psi(xy) = \psi(y)\psi(x)$

# $(\pm\beta)$ -integers $\cdots$ encoding by infinite word

Let  $\beta > 1$  be a Pisot number (algebraic integer, conjugates  $|\cdot| < 1$ ). Then:

$\mathbb{Z}_\beta^+$  : is an infinite set ,  
contains distances of lengths  $\Delta_0, \dots, \Delta_k$ , all  $\leq 1$  ,  
can be encoded by infinite word  $\mathbf{u}_\beta \in \{0, \dots, k\}^\mathbb{N}$  ,  
there exists a morphism  $\varphi$  fixing  $\mathbf{u}_\beta$  .

$\mathbb{Z}_{-\beta}$  : is an infinite set  $\Leftrightarrow \beta \geq \tau = (1 + \sqrt{5})/2$  ,  
contains distances of lengths  $\Delta'_0, \dots, \Delta'_{k'}$  ,  
can be encoded by biinfinite word  $\mathbf{v}_{-\beta} \in \{0, \dots, k'\}^\mathbb{Z}$  ,  
there exists an antimorphism  $\psi$  fixing  $\mathbf{v}_{-\beta}$  .

- $\varphi(xy) = \varphi(x)\varphi(y)$ ,  $\psi(xy) = \psi(y)\psi(x)$
- $\varphi^2, \psi^2 \cdots$  both morphisms, can be compared



# $(\pm\beta)$ -integers $\cdots$ encoding by infinite word

Let  $\beta > 1$  be a Pisot number (algebraic integer, conjugates  $|\cdot| < 1$ ). Then:

$\mathbb{Z}_\beta^+$  : is an infinite set ,

contains distances of lengths  $\Delta_0, \dots, \Delta_k$ , all  $\leq 1$ ,  
can be encoded by infinite word  $\mathbf{u}_\beta \in \{0, \dots, k\}^\mathbb{N}$ ,  
there exists a morphism  $\varphi$  fixing  $\mathbf{u}_\beta$ .

$\mathbb{Z}_{-\beta}$  : is an infinite set  $\Leftrightarrow \beta \geq \tau = (1 + \sqrt{5})/2$ ,

contains distances of lengths  $\Delta'_0, \dots, \Delta'_{k'}$ ,


can be encoded by biinfinite word  $\mathbf{v}_{-\beta} \in \{0, \dots, k'\}^\mathbb{Z}$ ,  
there exists an antimorphism  $\psi$  fixing  $\mathbf{v}_{-\beta}$ .

- $\varphi(xy) = \varphi(x)\varphi(y)$ ,  $\psi(xy) = \psi(y)\psi(x)$
- $\varphi^2, \psi^2 \cdots$  both morphisms, can be compared
- $\beta$  not Pisot  $\Rightarrow$  possibly infinite number of types of  $\Delta_i, \Delta'_i$

# $(\pm\beta)$ -integers $\cdots$ examples

Let  $\beta = \tau$ , the Golden ratio.


## Example $(\mathbb{Z}_\tau^+)$



# $(\pm\beta)$ -integers $\cdots$ examples

Let  $\beta = \tau$ , the Golden ratio.

## Example $(\mathbb{Z}_\tau^+)$




0

# $(\pm\beta)$ -integers $\cdots$ examples

Let  $\beta = \tau$ , the Golden ratio.

## Example $(\mathbb{Z}_\tau^+)$




$$0 \rightarrow 0\mathbf{1}$$

# $(\pm\beta)$ -integers $\cdots$ examples

Let  $\beta = \tau$ , the Golden ratio.

## Example $(\mathbb{Z}_\tau^+)$




$0 \rightarrow 01 \rightarrow 010$

# $(\pm\beta)$ -integers $\cdots$ examples

Let  $\beta = \tau$ , the Golden ratio.

## Example $(\mathbb{Z}_\tau^+)$




$0 \rightarrow 01 \rightarrow 010 \rightarrow 010\mathbf{01}$

# $(\pm\beta)$ -integers $\cdots$ examples

Let  $\beta = \tau$ , the Golden ratio.


## Example $(\mathbb{Z}_\tau^+)$



$0 \rightarrow 01 \rightarrow 010 \rightarrow 01001 \rightarrow 01001010$

$(\pm\beta)$ -integers  $\cdots$  examples

Let  $\beta = \tau$ , the Golden ratio.

Example  $(\mathbb{Z}_\tau^+)$ 


$0 \rightarrow 01 \rightarrow 010 \rightarrow 01001 \rightarrow 01001010 \rightarrow 01001010\mathbf{01001} \rightarrow \cdots$



# $(\pm\beta)$ -integers $\cdots$ examples


Let  $\beta = \tau$ , the Golden ratio.

## Example $(\mathbb{Z}_\tau^+)$



$0 \rightarrow 01 \rightarrow 010 \rightarrow 01001 \rightarrow 01001010 \rightarrow 0100101001001 \rightarrow \cdots$


## Example $(\mathbb{Z}_{-\tau})$



# $(\pm\beta)$ -integers $\cdots$ examples


Let  $\beta = \tau$ , the Golden ratio.

## Example $(\mathbb{Z}_\tau^+)$



$0 \rightarrow 01 \rightarrow 010 \rightarrow 01001 \rightarrow 01001010 \rightarrow 0100101001001 \rightarrow \cdots$

## Example $(\mathbb{Z}_{-\tau})$




1|0

# $(\pm\beta)$ -integers $\cdots$ examples


Let  $\beta = \tau$ , the Golden ratio.

## Example $(\mathbb{Z}_\tau^+)$



$0 \rightarrow 01 \rightarrow 010 \rightarrow 01001 \rightarrow 01001010 \rightarrow 0100101001001 \rightarrow \cdots$

## Example $(\mathbb{Z}_{-\tau})$




$1|0 \rightarrow 01|0$

# $(\pm\beta)$ -integers $\cdots$ examples


Let  $\beta = \tau$ , the Golden ratio.

## Example $(\mathbb{Z}_\tau^+)$



$0 \rightarrow 01 \rightarrow 010 \rightarrow 01001 \rightarrow 01001010 \rightarrow 0100101001001 \rightarrow \cdots$

## Example $(\mathbb{Z}_{-\tau})$




$1|0 \rightarrow 01|0 \rightarrow 01|001$

# $(\pm\beta)$ -integers $\cdots$ examples


Let  $\beta = \tau$ , the Golden ratio.

## Example $(\mathbb{Z}_\tau^+)$



$0 \rightarrow 01 \rightarrow 010 \rightarrow 01001 \rightarrow 01001010 \rightarrow 0100101001001 \rightarrow \cdots$

## Example $(\mathbb{Z}_{-\tau})$




$1|0 \rightarrow 01|0 \rightarrow 01|001 \rightarrow \textcolor{red}{001}01|001$

# $(\pm\beta)$ -integers $\cdots$ examples


Let  $\beta = \tau$ , the Golden ratio.

## Example $(\mathbb{Z}_\tau^+)$



$0 \rightarrow 01 \rightarrow 010 \rightarrow 01001 \rightarrow 01001010 \rightarrow 0100101001001 \rightarrow \cdots$

## Example $(\mathbb{Z}_{-\tau})$



$1|0 \rightarrow 01|0 \rightarrow 01|001 \rightarrow 00101|001 \rightarrow 00101|001\mathbf{00101} \rightarrow \cdots$

# Table of contents

1  $(\pm\beta)$ -integers

2 Similarity of  $\mathbb{Z}_\beta^+$  and  $\mathbb{Z}_{-\beta}$

3 Cubic Pisot examples

# Confluent Pisot numbers

For  $\beta$ -expansions, we have the following theorem.

**Theorem (Ch. Frougny)**

*Let  $\beta > 1$ . Then the following conditions are equivalent:*



# Confluent Pisot numbers

For  $\beta$ -expansions, we have the following theorem.

## Theorem (Ch. Frougny)

*Let  $\beta > 1$ . Then the following conditions are equivalent:*

①  $\mathbb{Z}_\beta^+ = \left\{ \sum_{i=0}^k a_i \beta^i : k \in \mathbb{N}, a_i \in \{0, 1, \dots, \lceil \beta \rceil - 1\} \right\}.$

# Confluent Pisot numbers

For  $\beta$ -expansions, we have the following theorem.

## Theorem (Ch. Frougny)

*Let  $\beta > 1$ . Then the following conditions are equivalent:*

- ①  $\mathbb{Z}_\beta^+ = \left\{ \sum_{i=0}^k a_i \beta^i : k \in \mathbb{N}, a_i \in \{0, 1, \dots, \lceil \beta \rceil - 1\} \right\}$ .
- ②  $\beta$  is a Pisot number with minimal polynomial  $x^d - mx^{d-1} - \dots - mx^2 - mx - n$  with  $m \geq n \geq 1$ .

# Confluent Pisot numbers

For  $\beta$ -expansions, we have the following theorem.

## Theorem (Ch. Frougny)

*Let  $\beta > 1$ . Then the following conditions are equivalent:*

- ①  $\mathbb{Z}_\beta^+ = \left\{ \sum_{i=0}^k a_i \beta^i : k \in \mathbb{N}, a_i \in \{0, 1, \dots, \lceil \beta \rceil - 1\} \right\}$ .
- ②  $\beta$  is a Pisot number with minimal polynomial  $x^d - mx^{d-1} - \dots - mx^2 - mx - n$  with  $m \geq n \geq 1$ .

Note:

- if  $z = \sum_{i=0}^k a_i \beta^i$ , then  $z \in \mathbb{Z}_\beta^+$ , even if  $a_k \cdots a_0 0^\omega$  is not admissible

# Confluent Pisot numbers

For  $\beta$ -expansions, we have the following theorem.

## Theorem (Ch. Frougny)

*Let  $\beta > 1$ . Then the following conditions are equivalent:*

- ①  $\mathbb{Z}_\beta^+ = \left\{ \sum_{i=0}^k a_i \beta^i : k \in \mathbb{N}, a_i \in \{0, 1, \dots, \lceil \beta \rceil - 1\} \right\}$ .
- ②  $\beta$  is a Pisot number with minimal polynomial  $x^d - mx^{d-1} - \dots - mx^2 - mx - n$  with  $m \geq n \geq 1$ .

Note:

- if  $z = \sum_{i=0}^k a_i \beta^i$ , then  $z \in \mathbb{Z}_\beta^+$ , even if  $a_k \cdots a_0 0^\omega$  is not admissible
- this class of  $\beta \cdots$  confluent Pisot numbers

# Similarity of $\mathbb{Z}_\beta^+$ and $\mathbb{Z}_{-\beta}$

For whose  $\beta$  are  $\mathbb{Z}_\beta^+$  and  $\mathbb{Z}_{-\beta}$  “similar”?

What is “similar”?

# Similarity of $\mathbb{Z}_\beta^+$ and $\mathbb{Z}_{-\beta}$

For whose  $\beta$  are  $\mathbb{Z}_\beta^+$  and  $\mathbb{Z}_{-\beta}$  “similar”?

What is “similar”?

- distances in  $\mathbb{Z}_{-\beta}$  are also at most 1

# Similarity of $\mathbb{Z}_\beta^+$ and $\mathbb{Z}_{-\beta}$

For whose  $\beta$  are  $\mathbb{Z}_\beta^+$  and  $\mathbb{Z}_{-\beta}$  “similar”?

What is “similar”?

- distances in  $\mathbb{Z}_{-\beta}$  are also at most 1
- the sets of distances in  $\mathbb{Z}_\beta^+$  and  $\mathbb{Z}_{-\beta}$  are exactly the same

# Similarity of $\mathbb{Z}_\beta^+$ and $\mathbb{Z}_{-\beta}$

For whose  $\beta$  are  $\mathbb{Z}_\beta^+$  and  $\mathbb{Z}_{-\beta}$  “similar”?

What is “similar”?

- distances in  $\mathbb{Z}_{-\beta}$  are also at most 1
- the sets of distances in  $\mathbb{Z}_\beta^+$  and  $\mathbb{Z}_{-\beta}$  are exactly the same
- the encodings  $\mathbf{u}_\beta$  and  $\mathbf{v}_{-\beta}$  have the same language



# Similarity of $\mathbb{Z}_\beta^+$ and $\mathbb{Z}_{-\beta}$

For whose  $\beta$  are  $\mathbb{Z}_\beta^+$  and  $\mathbb{Z}_{-\beta}$  “similar”?

What is “similar”?

- distances in  $\mathbb{Z}_{-\beta}$  are also at most 1
- the sets of distances in  $\mathbb{Z}_\beta^+$  and  $\mathbb{Z}_{-\beta}$  are exactly the same
- the encodings  $\mathbf{u}_\beta$  and  $\mathbf{v}_{-\beta}$  have the same language

Note:

- two infinite words have the same language iff morphisms fixing them are conjugated

# Conjugacy of morphisms

## Definition

*Let  $\mathcal{A}$  be an alphabet (finite or infinite) and  $\pi, \rho : \mathcal{A}^* \rightarrow \mathcal{A}^*$  be morphisms on  $\mathcal{A}$ . We say that  $\pi$  and  $\rho$  are conjugated, if there exists a word  $w \in \mathcal{A}^*$  such that either*

*$w\pi(a) = \rho(a)w$ , for all  $a \in \mathcal{A}$ , or  $\pi(a)w = w\rho(a)$ , for all  $a \in \mathcal{A}$ .*

*We denote  $\pi \sim \rho$ .*

# Conjugacy of morphisms

## Definition

Let  $\mathcal{A}$  be an alphabet (finite or infinite) and  $\pi, \rho : \mathcal{A}^* \rightarrow \mathcal{A}^*$  be morphisms on  $\mathcal{A}$ . We say that  $\pi$  and  $\rho$  are conjugated, if there exists a word  $w \in \mathcal{A}^*$  such that either

$$w\pi(a) = \rho(a)w, \text{ for all } a \in \mathcal{A}, \text{ or } \pi(a)w = w\rho(a), \text{ for all } a \in \mathcal{A}.$$

We denote  $\pi \sim \rho$ .

## Example

Let  $\beta = \tau$ ,  $\mathcal{A} = \{0, 1\}$ .

- $\mathbf{u}_\tau$  fixed point of  $\varphi^2$ :  $\varphi^2(0) = 010, \varphi^2(1) = 01$
- $\mathbf{v}_{-\tau}$  fixed point of  $\psi^2$ :  $\psi^2(0) = 001, \psi^2(1) = 01$
- $\varphi^2(a)(01) = (01)\psi^2(a)$

# Main result

Theorem (D., Z. Masáková, T. Vávra)

*Let  $\beta > 1$ . Then the following conditions are equivalent:*

# Main result

## Theorem (D., Z. Masáková, T. Vávra)

*Let  $\beta > 1$ . Then the following conditions are equivalent:*

- 1 Words  $\mathbf{u}_\beta$  and  $\mathbf{v}_{-\beta}$  have the same language ( $\varphi^2 \sim \psi^2$ ).

# Main result

## Theorem (D., Z. Masáková, T. Vávra)

*Let  $\beta > 1$ . Then the following conditions are equivalent:*

- 1 Words  $\mathbf{u}_\beta$  and  $\mathbf{v}_{-\beta}$  have the same language ( $\varphi^2 \sim \psi^2$ ).
- 2  $\mathbb{Z}_{-\beta} = \left\{ \sum_{i=0}^k a_i (-\beta)^i : k \in \mathbb{N}, a_j \in \{0, 1, \dots, \lfloor \beta \rfloor\} \right\}$ .

# Main result

## Theorem (D., Z. Masáková, T. Vávra)

*Let  $\beta > 1$ . Then the following conditions are equivalent:*

- ❶ *Words  $\mathbf{u}_\beta$  and  $\mathbf{v}_{-\beta}$  have the same language ( $\varphi^2 \sim \psi^2$ ).*
- ❷  *$\mathbb{Z}_{-\beta} = \left\{ \sum_{i=0}^k a_i(-\beta)^i : k \in \mathbb{N}, a_j \in \{0, 1, \dots, \lfloor \beta \rfloor\} \right\}$ .*
- ❸  *$\beta$  is a Pisot number with minimal polynomial  $x^d - mx^{d-1} - \dots - mx^2 - mx - n$  with  $m \geq n \geq 1$  for  $d$  odd and  $m = n \geq 1$  for  $d$  even.*

## Even degree remark

Let  $\beta > 1$  be a Pisot number with minimal polynomial

$x^d - mx^{d-1} - \dots - mx^2 - mx - n$  with  $d$  even and  $m > n \geq 1$ .



## Even degree remark

Let  $\beta > 1$  be a Pisot number with minimal polynomial  $x^d - mx^{d-1} - \dots - mx^2 - mx - n$  with  $d$  even and  $m > n \geq 1$ .

- $\Delta'_i = \Delta_i$  for  $i \leq k-2$ , but  $\Delta'_{k-1} = \Delta_{k-1} + 1$

## Even degree remark

Let  $\beta > 1$  be a Pisot number with minimal polynomial  $x^d - mx^{d-1} - \dots - mx^2 - mx - n$  with  $d$  even and  $m > n \geq 1$ .

- $\Delta'_i = \Delta_i$  for  $i \leq k-2$ , but  $\Delta'_{k-1} = \Delta_{k-1} + 1$
- we can cut  $\Delta'_{k-1} = 1 + \frac{n}{\beta} = \Delta_0 + \Delta_{k-1}$

# Even degree remark

Let  $\beta > 1$  be a Pisot number with minimal polynomial  $x^d - mx^{d-1} - \dots - mx^2 - mx - n$  with  $d$  even and  $m > n \geq 1$ .

- $\Delta'_i = \Delta_i$  for  $i \leq k-2$ , but  $\Delta'_{k-1} = \Delta_{k-1} + 1$
- we can cut  $\Delta'_{k-1} = 1 + \frac{n}{\beta} = \Delta_0 + \Delta_{k-1}$
- equivalent to application of morphism  $\pi$  on  $\mathbf{v}_{-\beta}$ , where

$$\pi(i) = \begin{cases} i & \text{if } i \in \{0, \dots, k-2\}, \\ 0(k-1) & \text{if } i = k-1. \end{cases}$$

## Even degree remark

Let  $\beta > 1$  be a Pisot number with minimal polynomial  $x^d - mx^{d-1} - \dots - mx^2 - mx - n$  with  $d$  even and  $m > n \geq 1$ .

- $\Delta'_i = \Delta_i$  for  $i \leq k-2$ , but  $\Delta'_{k-1} = \Delta_{k-1} + 1$
- we can cut  $\Delta'_{k-1} = 1 + \frac{n}{\beta} = \Delta_0 + \Delta_{k-1}$
- equivalent to application of morphism  $\pi$  on  $\mathbf{v}_{-\beta}$ , where

$$\pi(i) = \begin{cases} i & \text{if } i \in \{0, \dots, k-2\}, \\ 0(k-1) & \text{if } i = k-1. \end{cases}$$

- there exists an antimorphism  $\psi'$  with  $\pi(\mathbf{v}_{-\beta})$  as its fixed point

# Even degree remark

Let  $\beta > 1$  be a Pisot number with minimal polynomial  $x^d - mx^{d-1} - \dots - mx^2 - mx - n$  with  $d$  even and  $m > n \geq 1$ .

- $\Delta'_i = \Delta_i$  for  $i \leq k-2$ , but  $\Delta'_{k-1} = \Delta_{k-1} + 1$
- we can cut  $\Delta'_{k-1} = 1 + \frac{n}{\beta} = \Delta_0 + \Delta_{k-1}$
- equivalent to application of morphism  $\pi$  on  $\mathbf{v}_{-\beta}$ , where

$$\pi(i) = \begin{cases} i & \text{if } i \in \{0, \dots, k-2\}, \\ 0(k-1) & \text{if } i = k-1. \end{cases}$$

- there exists an antimorphism  $\psi'$  with  $\pi(\mathbf{v}_{-\beta})$  as its fixed point
- $\varphi^2 \sim \psi'^2$

# Table of contents

- 1  $(\pm\beta)$ -integers
- 2 Similarity of  $\mathbb{Z}_\beta^+$  and  $\mathbb{Z}_{-\beta}$
- 3 Cubic Pisot examples

# Cubic Pisot unit case

## Proposition

*Let  $\beta > 1$  be cubic Pisot unit with minimal polynomial*  
$$p(x) = x^3 - ax^2 - bx \pm 1.$$

# Cubic Pisot unit case

## Proposition

Let  $\beta > 1$  be cubic Pisot unit with minimal polynomial  
 $p(x) = x^3 - ax^2 - bx \pm 1$ .

- ① For  $p(x) = x^3 - mx^2 - mx - 1$ :  
the distances in  $\mathbb{Z}_\beta^+$  and  $\mathbb{Z}_{-\beta}$  are the same and  $\varphi_\beta^2 \sim \psi_{-\beta}^2$ .



# Cubic Pisot unit case

## Proposition

Let  $\beta > 1$  be cubic Pisot unit with minimal polynomial  $p(x) = x^3 - ax^2 - bx \pm 1$ .

- ① For  $p(x) = x^3 - mx^2 - mx - 1$ :  
the distances in  $\mathbb{Z}_\beta^+$  and  $\mathbb{Z}_{-\beta}$  are the same and  $\varphi_\beta^2 \sim \psi_{-\beta}^2$ .
- ② For  $p(x) = x^3 - mx^2 + x - 1$  and  $p(x) = x^3 - mx^2 + 1$ :  
the distances in  $\mathbb{Z}_\beta^+$  and  $\mathbb{Z}_{-\beta}$  are the same, but  $\varphi_\beta^2 \approx \psi_{-\beta}^2$ .

# Cubic Pisot unit case

## Proposition

Let  $\beta > 1$  be cubic Pisot unit with minimal polynomial  $p(x) = x^3 - ax^2 - bx \pm 1$ .

- ① For  $p(x) = x^3 - mx^2 - mx - 1$ :  
the distances in  $\mathbb{Z}_\beta^+$  and  $\mathbb{Z}_{-\beta}$  are the same and  $\varphi_\beta^2 \sim \psi_{-\beta}^2$ .
- ② For  $p(x) = x^3 - mx^2 + x - 1$  and  $p(x) = x^3 - mx^2 + 1$ :  
the distances in  $\mathbb{Z}_\beta^+$  and  $\mathbb{Z}_{-\beta}$  are the same, but  $\varphi_\beta^2 \approx \psi_{-\beta}^2$ .
- ③ In all other cases,  $\mathbb{Z}_{-\beta}$  contains distances  $> 1$ .

# Projection of $(\pm\beta)$ -integers

Let  $\beta > 1$  be cubic Pisot unit with conjugates  $\beta', \beta''$ .

## Projection of $(\pm\beta)$ -integers

Let  $\beta > 1$  be cubic Pisot unit with conjugates  $\beta', \beta''$ .

- $\mathbb{Z}_{\pm\beta}^{(+)} \subset \mathbb{R}$  is either single-element or unbounded aperiodic set

# Projection of ( $\pm\beta$ )-integers

Let  $\beta > 1$  be cubic Pisot unit with conjugates  $\beta', \beta''$ .

- $\mathbb{Z}_{\pm\beta}^{(+)} \subset \mathbb{R}$  is either single-element or unbounded aperiodic set
- possible projection to a bounded set in  $\mathbb{R}^2$  or  $\mathbb{C}$ 
  - for  $\beta', \beta'' \in \mathbb{R}$ :

$$\underbrace{\sum_{i=0}^k a_i (\pm\beta)^i}_{\in \mathbb{Z}_{\pm\beta}^{(+)}} \longrightarrow \underbrace{\left( \sum_{i=0}^k a_i (\pm\beta')^i, \sum_{i=0}^k a_i (\pm\beta'')^i \right)}_{\in \mathbb{R}^2}$$

# Projection of ( $\pm\beta$ )-integers

Let  $\beta > 1$  be cubic Pisot unit with conjugates  $\beta', \beta''$ .

- $\mathbb{Z}_{\pm\beta}^{(+)} \subset \mathbb{R}$  is either single-element or unbounded aperiodic set
- possible projection to a bounded set in  $\mathbb{R}^2$  or  $\mathbb{C}$ 
  - for  $\beta', \beta'' \in \mathbb{R}$ :

$$\underbrace{\sum_{i=0}^k a_i (\pm\beta)^i}_{\in \mathbb{Z}_{\pm\beta}^{(+)}} \longrightarrow \underbrace{\left( \sum_{i=0}^k a_i (\pm\beta')^i, \sum_{i=0}^k a_i (\pm\beta'')^i \right)}_{\in \mathbb{R}^2}$$

- for  $\beta', \beta'' \in \mathbb{C} \setminus \mathbb{R}$ :


$$\underbrace{\sum_{i=0}^k a_i (\pm\beta)^i}_{\in \mathbb{Z}_{\pm\beta}^{(+)}} \longrightarrow \underbrace{\sum_{i=0}^k a_i (\pm\beta')^i}_{\in \mathbb{C}}$$

- what we get  $\rightarrow$  Rauzy fractal


Case 1: same distances,  $\varphi^2 \sim \psi^2$

# Case 1: same distances, $\varphi^2 \sim \psi^2$

$$p(x) = x^3 - x^2 - x - 1$$



$\mathbb{Z}_\beta^+$





$\mathbb{Z}_{-\beta}$





# Case 1: same distances, $\varphi^2 \sim \psi^2$

$$p(x) = x^3 - x^2 - x - 1$$




$$p(x) = x^3 - 3x^2 - 3x - 1$$




Case 2: same distances,  $\varphi^2 \approx \psi^2$

## Case 2: same distances, $\varphi^2 \approx \psi^2$

$$p(x) = x^3 - 2x^2 + x - 1$$





$\mathbb{Z}_\beta^+$





$\mathbb{Z}_{-\beta}$

## Case 2: same distances, $\varphi^2 \approx \psi^2$

$$p(x) = x^3 - 2x^2 + x - 1$$




$$p(x) = x^3 - 3x^2 + 1$$




## Case 3: different distances

## Case 3: different distances

$$p(x) = x^3 - x^2 - 2x - 1$$




$\mathbb{Z}_\beta^+$



$\mathbb{Z}_{-\beta}$


# Case 3: different distances

$$p(x) = x^3 - x^2 - 2x - 1$$




$\mathbb{Z}_\beta^+$


$$p(x) = x^3 - 3x^2 + 2x - 1$$



$\mathbb{Z}_\beta^+$



$\mathbb{Z}_{-\beta}$



$\mathbb{Z}_{-\beta}$