

The Lexicographic Cross-Section of the Plactic Monoid is Regular

Robert Mercas

Christian-Albrechts-Universität zu Kiel



WORDS 2013 - Turku

A joint work with Christian Choffrut (LIAFA)

Schensted's Algorithm

QUESTION

Given a sequence, how long is its longest nondecreasing subword?

Schensted's Algorithm

QUESTION

Given a sequence, how long is its longest nondecreasing subword?

132541

Schensted's Algorithm

QUESTION

Given a sequence, how long is its longest nondecreasing subword?

132541

125

Schensted's Algorithm

QUESTION

Given a sequence, how long is its longest nondecreasing subword?

132541

124

Schensted's Algorithm

QUESTION

Given a sequence, how long is its longest nondecreasing subword?

132541

135

Schensted's Algorithm

QUESTION

Given a sequence, how long is its longest nondecreasing subword?

132541

134

Schensted's Algorithm

QUESTION

Given a sequence, how long is its longest nondecreasing subword?

132541

125, 124, 135, 134

1
←

Schensted's Algorithm

QUESTION

Given a sequence, how long is its longest nondecreasing subword?

132541

125, 124, 135, 134

1

3

Schensted's Algorithm

QUESTION

Given a sequence, how long is its longest nondecreasing subword?

132541

125, 124, 135, 134

1 3

2

Schensted's Algorithm

QUESTION

Given a sequence, how long is its longest nondecreasing subword?

132541

125, 124, 135, 134

1 2

3
←

Schensted's Algorithm

QUESTION

Given a sequence, how long is its longest nondecreasing subword?

132541

125, 124, 135, 134




Schensted's Algorithm

QUESTION

Given a sequence, how long is its longest nondecreasing subword?

132541

125, 124, 135, 134




Schensted's Algorithm

QUESTION

Given a sequence, how long is its longest nondecreasing subword?

132541

125, 124, 135, 134




Schensted's Algorithm

QUESTION

Given a sequence, how long is its longest nondecreasing subword?

132541

125, 124, 135, 134




Schensted's Algorithm

QUESTION

Given a sequence, how long is its longest nondecreasing subword?

132541

125, 124, 135, 134



2


Schensted's Algorithm

QUESTION

Given a sequence, how long is its longest nondecreasing subword?

132541

125, 124, 135, 134



3
←


Schensted's Algorithm

QUESTION

Given a sequence, how long is its longest nondecreasing subword?

132541

125, 124, 135, 134



Definition: Young Tableaux

- ▶ **Row:** nondecreasing sequence;

Definition: Young Tableaux

- ▶ **Row**: nondecreasing sequence;
- ▶ **Column**: strictly decreasing sequence;

Definition: Young Tableaux

- ▶ **Row**: nondecreasing sequence;
- ▶ Row u *dominates* row v , denoted $u \succeq v$ if $|u| \leq |v|$ and $u_i \geq v_i$, where $i = 1, \dots, |v|$.
- ▶ **Column**: strictly decreasing sequence;
- ▶ Column u *dominates* column v , denoted $u \succeq v$ if $|u| \geq |v|$ and $u_{|u|-|v|+i} \geq v_i$, where $i = 1, \dots, |v|$.

Definition: Young Tableaux

- ▶ **Row**: nondecreasing sequence;
- ▶ Row u *dominates* row v , denoted $u \succeq v$ if $|u| \leq |v|$ and $u_i \geq v_i$, where $i = 1, \dots, |v|$.
- ▶ **Column**: strictly decreasing sequence;
- ▶ Column u *dominates* column v , denoted $u \succeq v$ if $|u| \geq |v|$ and $u_{|u|-|v|+i} \geq v_i$, where $i = 1, \dots, |v|$.
- ▶ **Young tableau**: graphical representation of a sequence of columns $v_1 \succeq \dots \succeq v_p$.

Definition: Young Tableaux

- ▶ **Row**: nondecreasing sequence;
- ▶ Row u *dominates* row v , denoted $u \succeq v$ if $|u| \leq |v|$ and $u_i \geq v_i$, where $i = 1, \dots, |v|$.
- ▶ **Column**: strictly decreasing sequence;
- ▶ Column u *dominates* column v , denoted $u \succeq v$ if $|u| \geq |v|$ and $u_{|u|-|v|+i} \geq v_i$, where $i = 1, \dots, |v|$.
- ▶ **Young tableau**: graphical representation of a sequence of columns $v_1 \succeq \dots \succeq v_p$.

132541

Definition: Young Tableaux

- ▶ **Row**: nondecreasing sequence;
- ▶ Row u *dominates* row v , denoted $u \succeq v$ if $|u| \leq |v|$ and $u_i \geq v_i$, where $i = 1, \dots, |v|$.
- ▶ **Column**: strictly decreasing sequence;
- ▶ Column u *dominates* column v , denoted $u \succeq v$ if $|u| \geq |v|$ and $u_{|u|-|v|+i} \geq v_i$, where $i = 1, \dots, |v|$.
- ▶ **Young tableau**: graphical representation of a sequence of columns $v_1 \succeq \dots \succeq v_p$.


132541

3		
2	5	
1	1	4

Definition: Young Tableaux

- ▶ **Row**: nondecreasing sequence;
- ▶ Row u *dominates* row v , denoted $u \succeq v$ if $|u| \leq |v|$ and $u_i \geq v_i$, where $i = 1, \dots, |v|$.
- ▶ **Column**: strictly decreasing sequence;
- ▶ Column u *dominates* column v , denoted $u \succeq v$ if $|u| \geq |v|$ and $u_{|u|-|v|+i} \geq v_i$, where $i = 1, \dots, |v|$.
- ▶ **Young tableau**: graphical representation of a sequence of columns $v_1 \succeq \dots \succeq v_p$.

132541



321 51 4

3 25 114

- ▶ *monoïde plaxique* [Lascoux, Schtzenberger, '81]

- ▶ *monoïde plaxique* [Lascoux, Schtzenberger, '81] - it may refer to plate tectonics (tectonique des plaques in French), as the action of a generator of the plactic monoid resembles plates sliding past each other in an earthquake.

- ▶ *monoïde plaxique* [Lascoux, Schtzenberger, '81] - it may refer to plate tectonics (tectonique des plaques in French), as the action of a generator of the plactic monoid resembles plates sliding past each other in an earthquake.
- ▶ No results for either *plactic* or *plaxic*

- ▶ *monoïde plaxique* [Lascoux, Schtzenberger, '81] - it may refer to plate tectonics (tectonique des plaques in French), as the action of a generator of the plactic monoid resembles plates sliding past each other in an earthquake.
- ▶ No results for either *plactic* or *plaxic*
Latin *plaxus* under the etymology for the obsolete word *plash* (To bend down and interweave (stems partly cut through, branches, and twigs) so as to form a hedge or fence.): an unattested post-classical Latin form *plaxus*, alteration of classical Latin *plexus*, past participle of *plectere* (to plait, interweave, twine)

Definition: plactic monoid

- ▶ Take free monoid Σ^*

Definition: plactic monoid

- ▶ Take free monoid Σ^*
- ▶ Take the following relation R :

$$bac \equiv bca, \text{ where } a < b \leq c$$

$$acb \equiv cab, \text{ where } a \leq b < c$$

Definition: plactic monoid

► Take free monoid Σ^*

► Take the following relation R :

$$bac \equiv bca, \text{ where } a < b \leq c$$

$$acb \equiv cab, \text{ where } a \leq b < c$$

► **Plactic monoid**: the quotient of Σ^* by the congruence generated by R

Definition: plactic monoid

- ▶ Take free monoid Σ^*
- ▶ Take the following relation R :
$$bac \equiv bca, \text{ where } a < b \leq c$$
$$acb \equiv cab, \text{ where } a \leq b < c$$
- ▶ **Plactic monoid**: the quotient of Σ^* by the congruence generated by R

132541

Definition: plactic monoid

► Take free monoid Σ^*

► Take the following relation R :

$$bac \equiv bca, \text{ where } a < b \leq c$$

$$acb \equiv cab, \text{ where } a \leq b < c$$

► **Plactic monoid**: the quotient of Σ^* by the congruence generated by R

132541

\rightarrow 312541

Definition: plactic monoid

- ▶ Take free monoid Σ^*
- ▶ Take the following relation R :

$$bac \equiv bca, \text{ where } a < b \leq c$$

$$acb \equiv cab, \text{ where } a \leq b < c$$

- ▶ **Plactic monoid**: the quotient of Σ^* by the congruence generated by R

132541

$\rightarrow 312541 \rightarrow 315241$

Definition: plactic monoid

► Take free monoid Σ^*

► Take the following relation R :

$$bac \equiv bca, \text{ where } a < b \leq c$$

$$acb \equiv cab, \text{ where } a \leq b < c$$

► **Plactic monoid**: the quotient of Σ^* by the congruence generated by R

132541

$\rightarrow 312541 \rightarrow 315\textcolor{red}{24}1 \rightarrow 315214$

Definition: plactic monoid

- ▶ Take free monoid Σ^*
- ▶ Take the following relation R :

$$bac \equiv bca, \text{ where } a < b \leq c$$

$$acb \equiv cab, \text{ where } a \leq b < c$$

- ▶ **Plactic monoid**: the quotient of Σ^* by the congruence generated by R

132541

$\rightarrow 312541 \rightarrow 315241 \rightarrow 315214 \rightarrow 351214$

Definition: plactic monoid

► Take free monoid Σ^*

► Take the following relation R :

$$bac \equiv bca, \text{ where } a < b \leq c$$

$$acb \equiv cab, \text{ where } a \leq b < c$$

► **Plactic monoid**: the quotient of Σ^* by the congruence generated by R

132541

$\rightarrow 312541 \rightarrow 315241 \rightarrow 315214 \rightarrow 35\textcolor{red}{1}2\textcolor{red}{1}4 \rightarrow 352114$

Definition: plactic monoid

► Take free monoid Σ^*

► Take the following relation R :

$$bac \equiv bca, \text{ where } a < b \leq c$$

$$acb \equiv cab, \text{ where } a \leq b < c$$

► **Plactic monoid**: the quotient of Σ^* by the congruence generated by R

132541

$\rightarrow 312541 \rightarrow 315241 \rightarrow 315214 \rightarrow 351214 \rightarrow 352114 \rightarrow 325114$

Definition: plactic monoid

► Take free monoid Σ^*

► Take the following relation R :

$$bac \equiv bca, \text{ where } a < b \leq c$$

$$acb \equiv cab, \text{ where } a \leq b < c$$

► **Plactic monoid**: the quotient of Σ^* by the congruence generated by R

132541

$\rightarrow 312541 \rightarrow 315241 \rightarrow 315214 \rightarrow 351214 \rightarrow 352114 \rightarrow 32\textcolor{red}{5}1\textcolor{red}{1}4 \rightarrow$

321 51 4

1 Cross-sections

2 Further remarks

Definition: Cross-sections

- ▶ **Cross-section** of an equivalence relation is a set of elements such that each class has exactly one element in the cross-section.

Definition: Cross-sections

- ▶ **Cross-section** of an equivalence relation is a set of elements such that each class has exactly one element in the cross-section.
- ▶ Denote $Y(u)$ the Young tableau equivalent to $u \in \Sigma^*$.

Definition: Cross-sections

- ▶ **Cross-section** of an equivalence relation is a set of elements such that each class has exactly one element in the cross-section.
- ▶ Denote $Y(u)$ the Young tableau equivalent to $u \in \Sigma^*$.
The plactic monoid admits a regular cross-section.

Definition: Cross-sections

- ▶ **Cross-section** of an equivalence relation is a set of elements such that each class has exactly one element in the cross-section.
- ▶ Denote $Y(u)$ the Young tableau equivalent to $u \in \Sigma^*$.
The plactic monoid admits a regular cross-section.
- ▶ AIM: Cross-section of the lexicographical representatives

LEMMA (BILLIARD RULE FOR ROWS)

Let $u \in \Sigma^$ be a row and let $a < u[|u|]$. Then $ua \equiv bxay$, where $xby = u$ and b is the leftmost element strictly greater than a .*

Example:

LEMMA (BILLIARD RULE FOR ROWS)

Let $u \in \Sigma^$ be a row and let $a < u[|u|]$. Then $ua \equiv bxay$, where $xb y = u$ and b is the leftmost element strictly greater than a .*

Example:

122345 · 2



LEMMA (BILLIARD RULE FOR ROWS)

Let $u \in \Sigma^$ be a row and let $a < u[|u|]$. Then $ua \equiv bxay$, where $xb y = u$ and b is the leftmost element strictly greater than a .*

Example:

3 · 122245



LEMMA (BILLIARD RULE FOR COLUMNS)

Let $u \in \Sigma^$ be a column and let $a \leq u[1]$. Then $au \equiv xayb$, where $xby = u$ and b is the least element greater than or equal to a .*

Example:

Billiard Rules (cont)

LEMMA (BILLIARD RULE FOR COLUMNS)

Let $u \in \Sigma^$ be a column and let $a \leq u[1]$. Then $au \equiv xayb$, where $xby = u$ and b is the least element greater than or equal to a .*

Example:



$$3 \cdot 54\underline{3}21$$

Billiard Rules (cont)

LEMMA (BILLIARD RULE FOR COLUMNS)

Let $u \in \Sigma^$ be a column and let $a \leq u[1]$. Then $au \equiv xayb$, where $xby = u$ and b is the least element greater than or equal to a .*

Example:



54321 · 3

Billiard Rules (cont)

LEMMA (BILLIARD RULE FOR COLUMNS)

Let $u \in \Sigma^$ be a column and let $a \leq u[1]$. Then $au \equiv xayb$, where $xby = u$ and b is the least element greater than or equal to a .*

Example:



$$54\underline{3}21 \cdot 3$$



$$2 \cdot 54\underline{3}1$$

Billiard Rules (cont)

LEMMA (BILLIARD RULE FOR COLUMNS)

Let $u \in \Sigma^$ be a column and let $a \leq u[1]$. Then $au \equiv xayb$, where $xby = u$ and b is the least element greater than or equal to a .*

Example:



$$54\underline{3}21 \cdot 3$$



$$54\underline{2}1 \cdot 3$$

PROPOSITION

Let u and v be two columns defining a Young tableau, i.e., $u \succcurlyeq v$. Let w be a column such that $w[1] < u[|u|]$. Then $uvw \equiv uwv$.

PROPOSITION

Let u and v be two columns defining a Young tableau, i.e., $u \succcurlyeq v$. Let w be a column such that $w[1] < u[|u|]$. Then $uvw \equiv uwv$.

COROLLARY

Let u and v be two columns such that v is a subword of u . Then $uv \equiv vu$.

PROPOSITION

The maximum representative of the congruence class a word $v \in \{1, 2\}^$ is part of, is of the form*

$$v_{\max} = 2^m 1^{|v|_1} 2^{|v|_2 - m},$$

where $m < |v|_1$ or $|v|_2 - m = 0$.

Maximal cross-section

PROPOSITION

The maximum representative of the congruence class a word $v \in \{1, 2\}^$ is part of, is of the form*

$$v_{\max} = 2^m 1^{|v|_1} 2^{|v|_2 - m},$$

where $m < |v|_1$ or $|v|_2 - m = 0$.

Proof: [Arnold, Kanta, Krob, '97] $(21)^* 1^* 2^*$ is a cross-section.

For binary alphabets 21 commutes with both 1 and 2.

$2m$ is the maximal length Dyck subword in v .

Maximal cross-section

PROPOSITION

The maximum representative of the congruence class a word $v \in \{1, 2\}^$ is part of, is of the form*

$$v_{\max} = 2^m 1^{|v|_1} 2^{|v|_2 - m},$$

where $m < |v|_1$ or $|v|_2 - m = 0$.

Proof: [Arnold, Kanta, Krob, '97] $(21)^* 1^* 2^*$ is a cross-section.

For binary alphabets 21 commutes with both 1 and 2.

$2m$ is the maximal length Dyck subword in v .

THEOREM

The language determined by the maximum representatives of the congruence class determined by Young tableaux is not regular.

PROPOSITION

The minimum representative of the congruence class a word $v \in \{1, 2\}^$ is part of, is of the form $v_{min} = 1^*(21)^*2^*$.*

PROPOSITION

The minimum representative of the congruence class a word $v \in \{1, 2\}^$ is part of, is of the form $v_{min} = 1^*(21)^*2^*$.*

PROPOSITION

The minimum representative of the congruence class a word $v \in \{1, 2, 3\}^$ is part of, is of the form $v_{min} = 1^*(21)^*(2^* + (31)^*)(321)^*(32)^*3^*$.*

For $Y(u)$ we associate the *decomposition* $(n_1^{e_1}, \dots, n_p^{e_p})$ where e_1 are its first columns of same height n_1 , e_2 next columns of height n_2 , \dots , etc., and the sequence n_1, \dots, n_p is strictly decreasing.

For $Y(u)$ we associate the *decomposition* $(n_1^{e_1}, \dots, n_p^{e_p})$ where e_1 are its first columns of same height n_1 , e_2 next columns of height n_2 , \dots , etc., and the sequence n_1, \dots, n_p is strictly decreasing.

PROPOSITION

If $u = av$, for some $a \in \Sigma$, then $Y(v)$ has a decomposition of the form

$$(n_1^{e_1}, \dots, n_{i-1}^{e_{i-1}}, n_i^{e_i-1}, n_i - 1, n_{i+1}^{e_{i+1}}, \dots, n_p^{e_p})$$

if $n_i > n_{i+1} + 1$, or, otherwise,

$$(n_1^{e_1}, \dots, n_{i-1}^{e_{i-1}}, n_i^{e_i-1}, n_{i+1}^{e_{i+1}+1}, \dots, n_r^{e_r}).$$

For $Y(u)$ we associate the *decomposition* $(n_1^{e_1}, \dots, n_p^{e_p})$ where e_1 are its first columns of same height n_1 , e_2 next columns of height n_2 , \dots , etc., and the sequence n_1, \dots, n_p is strictly decreasing.

PROPOSITION

If $u = av$, for some $a \in \Sigma$, then $Y(v)$ has a decomposition of the form

$$(n_1^{e_1}, \dots, n_{i-1}^{e_{i-1}}, n_i^{e_i-1}, n_i - 1, n_{i+1}^{e_{i+1}}, \dots, n_p^{e_p})$$


if $n_i > n_{i+1} + 1$, or, otherwise,

$$(n_1^{e_1}, \dots, n_{i-1}^{e_{i-1}}, n_i^{e_i-1}, n_{i+1}^{e_{i+1}+1}, \dots, n_r^{e_r}).$$


COROLLARY

Let x be an element of the plactic monoid. There exist k factorizations of the form $x = ay$, where $a \in \Sigma$ and k is the number of different lengths for the columns of the associated Young tableau.

EXAMPLE




EXAMPLE



1 32 541


EXAMPLE



1 32 541

1
↙

EXAMPLE




1 32 541



3
←

EXAMPLE




1 32 541



2

EXAMPLE




1 32 541



3
←

EXAMPLE




1 32 541




5
←

EXAMPLE




1 32 541




4
←

EXAMPLE




1 32 541




5

EXAMPLE




1 32 541



1
←

EXAMPLE




1 32 541




$\xrightarrow{2}$

EXAMPLE




1 32 541




3
←


EXAMPLE





1 32 541




EXAMPLE





1 32 541




EXAMPLE




1 3 2 5 4 1





\rightarrow 1.




EXAMPLE




1 32 541




\rightarrow 1.





\rightarrow 13.




EXAMPLE




1 3 2 5 4 1



$\rightarrow 1 \cdot$




$\rightarrow 13 \cdot$





$\rightarrow 132 \cdot$



EXAMPLE




1 32 541




\rightarrow

1.




\rightarrow

13.




\rightarrow



132.



EXAMPLE




1 3 2 5 4 1




\rightarrow

1.




\rightarrow

13.




\rightarrow

132.




\rightarrow



1.



EXAMPLE



1 3 2 5 4 1




\rightarrow

1.




\rightarrow

13.




\rightarrow

132.




\rightarrow



2.



EXAMPLE



1 32 541




\rightarrow

1.




\rightarrow

13.




\rightarrow

132.




\rightarrow



2.



EXAMPLE




1 32 541




\rightarrow

1.




\rightarrow

13.




\rightarrow

132.



\rightarrow 2.



Relations on columns

We write $u \trianglelefteq v$ if the following conditions hold:

- ▶ for all $i = 1, \dots, \min\{|u|, |v|\}$, the condition $u_i \leq v_i$ holds;
- ▶ furthermore, if $|u| < |v|$, then $u_{|u|} \leq v_{|u|+1}$ holds.

Relations on columns

We write $u \trianglelefteq v$ if the following conditions hold:

- ▶ for all $i = 1, \dots, \min\{|u|, |v|\}$, the condition $u_i \leq v_i$ holds;
- ▶ furthermore, if $|u| < |v|$, then $u_{|u|} \leq v_{|u|+1}$ holds.

$$432 \trianglelefteq 43 \text{ and } 43 \trianglelefteq 6541$$

Relations on columns

We write $u \trianglelefteq v$ if the following conditions hold:

- ▶ for all $i = 1, \dots, \min\{|u|, |v|\}$, the condition $u_i \leq v_i$ holds;
- ▶ furthermore, if $|u| < |v|$, then $u_{|u|} \leq v_{|u|+1}$ holds.

$432 \trianglelefteq 43$ and $43 \trianglelefteq 6541$, but $432 \not\trianglelefteq 6541$.

Relations on columns

We write $u \trianglelefteq v$ if the following conditions hold:

- ▶ for all $i = 1, \dots, \min\{|u|, |v|\}$, the condition $u_i \leq v_i$ holds;
- ▶ furthermore, if $|u| < |v|$, then $u_{|u|} \leq v_{|u|+1}$ holds.

$432 \trianglelefteq 43$ and $43 \trianglelefteq 6541$, but $432 \not\trianglelefteq 6541$.

($432 \trianglelefteq 43$ and $43 \trianglelefteq 6541$ minimal, but not $432 \trianglelefteq 43 \trianglelefteq 6541 \equiv 4321 \trianglelefteq 43 \trianglelefteq 654$.)

Relations on columns

We write $u \trianglelefteq v$ if the following conditions hold:

- ▶ for all $i = 1, \dots, \min\{|u|, |v|\}$, the condition $u_i \leq v_i$ holds;
- ▶ furthermore, if $|u| < |v|$, then $u_{|u|} \leq v_{|u|+1}$ holds.

$432 \trianglelefteq 43$ and $43 \trianglelefteq 6541$, but $432 \not\trianglelefteq 6541$.

($432 \trianglelefteq 43$ and $43 \trianglelefteq 6541$ minimal, but not $432 \trianglelefteq 6541 \equiv 4321 \trianglelefteq 43 \trianglelefteq 654$.)

THEOREM

The transitive closure of \trianglelefteq is antisymmetric.

Relations on columns

We write $u \trianglelefteq v$ if the following conditions hold:

- ▶ for all $i = 1, \dots, \min\{|u|, |v|\}$, the condition $u_i \leq v_i$ holds;
- ▶ furthermore, if $|u| < |v|$, then $u_{|u|} \leq v_{|u|+1}$ holds.

$432 \trianglelefteq 43$ and $43 \trianglelefteq 6541$, but $432 \not\trianglelefteq 6541$.

($432 \trianglelefteq 43$ and $43 \trianglelefteq 6541$ minimal, but not $432 \trianglelefteq 43 \trianglelefteq 6541 \equiv 4321 \trianglelefteq 43 \trianglelefteq 654$.)

THEOREM

The transitive closure of \trianglelefteq is antisymmetric.

$$z_0 \trianglelefteq \dots \trianglelefteq z_p,$$

with $z_0 = z_p$, then for all $0 < i < p$, we have $z_i = z_0$.

Characterization of minimal cross-section

PROPOSITION

Let u, v be two columns such that, neither uv nor vu is a column. Then uv is lexicographically minimum in its class if and only if $u \trianglelefteq v$ holds.

Characterization of minimal cross-section

PROPOSITION

Let u, v be two columns such that, neither uv nor vu is a column. Then uv is lexicographically minimum in its class if and only if $u \trianglelefteq v$ holds.

THEOREM

Let $u = u_1 \cdots u_n$ be a factorization of maximal strictly decreasing strings. Then, u is the minimal lexicographical element of the cross-section if and only if $u_i \trianglelefteq u_j$, for all $1 \leq i < j \leq n$.

Characterization of minimal cross-section

PROPOSITION

Let u, v be two columns such that, neither uv nor vu is a column. Then uv is lexicographically minimum in its class if and only if $u \trianglelefteq v$ holds.

THEOREM

Let $u = u_1 \cdots u_n$ be a factorization of maximal strictly decreasing strings. Then, u is the minimal lexicographical element of the cross-section if and only if $u_i \trianglelefteq u_j$, for all $1 \leq i < j \leq n$.

9
4 8
3 6 7 6
2 5 6 1

Characterization of minimal cross-section

PROPOSITION

Let u, v be two columns such that, neither uv nor vu is a column. Then uv is lexicographically minimum in its class if and only if $u \trianglelefteq v$ holds.

THEOREM

Let $u = u_1 \cdots u_n$ be a factorization of maximal strictly decreasing strings. Then, u is the minimal lexicographical element of the cross-section if and only if $u_i \trianglelefteq u_j$, for all $1 \leq i < j \leq n$.

$$\begin{array}{cccc} & 9 & & 9 \\ 4 & 8 & & 6 \ 8 \\ 3 \ 6 \ 7 \ 6 & = & 4 \ 5 \ 7 \ 6 \\ \textcolor{red}{2} \ 5 \ 6 \ 1 & & 3 \ \textcolor{red}{2} \ 6 \ 1 \end{array}$$

Characterization of minimal cross-section

PROPOSITION

Let u, v be two columns such that, neither uv nor vu is a column. Then uv is lexicographically minimum in its class if and only if $u \trianglelefteq v$ holds.

THEOREM

Let $u = u_1 \cdots u_n$ be a factorization of maximal strictly decreasing strings. Then, u is the minimal lexicographical element of the cross-section if and only if $u_i \trianglelefteq u_j$, for all $1 \leq i < j \leq n$.

$$\begin{array}{cccc} & 9 & & 9 & & 9 \\ 4 & 8 & & 6 & 8 & & 7 & 8 \\ 3 & 6 & 7 & 6 & = & 4 & 5 & 7 & 6 & = & 4 & 6 & 6 & 6 \\ 2 & 5 & 6 & 1 & & 3 & 2 & 6 & 1 & & 3 & 5 & 2 & 1 \end{array}$$

Characterization of minimal cross-section

PROPOSITION

Let u, v be two columns such that, neither uv nor vu is a column. Then uv is lexicographically minimum in its class if and only if $u \trianglelefteq v$ holds.

THEOREM

Let $u = u_1 \cdots u_n$ be a factorization of maximal strictly decreasing strings. Then, u is the minimal lexicographical element of the cross-section if and only if $u_i \trianglelefteq u_j$, for all $1 \leq i < j \leq n$.

$$\begin{array}{ccccccc} & 9 & & 9 & & 9 & & 7 \\ 4 & 8 & & 6 & 8 & & 7 & 8 & & 6 & 9 \\ 3 & 6 & 7 & 6 & = & 4 & 5 & 7 & 6 & = & 4 & 6 & 6 & 6 & = & 4 & 6 & 2 & 8 \\ 2 & 5 & 6 & 1 & & 3 & 2 & 6 & 1 & & 3 & 5 & 2 & \color{red}{1} & & 3 & 5 & \color{red}{1} & 6 \end{array}$$

Characterization of minimal cross-section

PROPOSITION

Let u, v be two columns such that, neither uv nor vu is a column. Then uv is lexicographically minimum in its class if and only if $u \trianglelefteq v$ holds.

THEOREM

Let $u = u_1 \cdots u_n$ be a factorization of maximal strictly decreasing strings. Then, u is the minimal lexicographical element of the cross-section if and only if $u_i \trianglelefteq u_j$, for all $1 \leq i < j \leq n$.

$$\begin{array}{ccccc} & 9 & & 9 & & 9 & & 7 & & 6 \\ 4 & 8 & & 6 & 8 & & 7 & 8 & & 6 & 9 \\ 3 & 6 & 7 & 6 & = & 4 & 5 & 7 & 6 & = & 4 & 6 & 6 & 6 & = & 4 & 6 & \textcolor{red}{2} & 8 & = & 4 & \textcolor{red}{2} & 7 & 8 \\ 2 & 5 & 6 & 1 & & 3 & 2 & 6 & 1 & & 3 & 5 & 2 & 1 & & 3 & 5 & \textcolor{red}{1} & 6 & & 3 & \textcolor{red}{1} & 6 & 6 \end{array}$$

Characterization of minimal cross-section

PROPOSITION

Let u, v be two columns such that, neither uv nor vu is a column. Then uv is lexicographically minimum in its class if and only if $u \trianglelefteq v$ holds.

THEOREM

Let $u = u_1 \cdots u_n$ be a factorization of maximal strictly decreasing strings. Then, u is the minimal lexicographical element of the cross-section if and only if $u_i \trianglelefteq u_j$, for all $1 \leq i < j \leq n$.

$$\begin{array}{cccccc}
 & 9 & & 9 & & 9 & & 7 & & 6 & & 4 \\
 4 & 8 & & 6 & 8 & & 7 & 8 & & 6 & 9 & & 5 & 9 & & 3 & 9 \\
 3 & 6 & 7 & 6 & = & 4 & 5 & 7 & 6 & = & 4 & 6 & 6 & 6 & = & 4 & 6 & 2 & 8 & = & 4 & 2 & 7 & 8 & = & 2 & 6 & 7 & 8 \\
 2 & 5 & 6 & 1 & & 3 & 2 & 6 & 1 & & 3 & 5 & 2 & 1 & & 3 & 5 & 1 & 6 & & 3 & 1 & 6 & 6 & & 1 & 5 & 6 & 6
 \end{array}$$


4
3 9
2 6 7 8
1 5 6 6

$$\begin{array}{cccc} 4 & & & 4\ 6\ 7\ 9 \\ 3 & & 9 & 3\ 5\ 6\ 8 \\ 2\ 6\ 7\ 8 & \rightarrow & 2 & 6 \\ 1\ 5\ 6\ 6 & & 1 & \end{array}$$

$$\begin{array}{ccccccc}
 4 & & & & 4 & 6 & 7 & 9 & & 4 & 6 & 7 & 9 \\
 3 & & 9 & & 3 & 5 & 6 & 8 & & 3 & 5 & 6 & 8 \\
 2 & 6 & 7 & 8 & \rightarrow & 2 & & 6 & \rightarrow & & & 2 & 6 \\
 1 & 5 & 6 & 6 & & 1 & & & & & & 1
 \end{array}$$

$$\begin{array}{ccccccc}
 4 & & & & 4 & 6 & 7 & 9 & & 4 & 6 & 7 & 9 & & \begin{array}{|c|c|c|c|} \hline 4 & 6 & 7 & 9 \\ \hline 3 & 5 & 6 & 8 \\ \hline & & 2 & 6 \\ \hline & & & 1 \\ \hline \end{array} \\
 3 & & 9 & & 3 & 5 & 6 & 8 & & 3 & 5 & 6 & 8 & = & \\
 2 & 6 & 7 & 8 & \rightarrow & 2 & & 6 & \rightarrow & & 2 & 6 & & & \\
 1 & 5 & 6 & 6 & & 1 & & & & & & 1 & & &
 \end{array}$$

contretableau



contretableau

COROLLARY

The minimal representative of a class has the same column length distribution as its Young tableau.

COROLLARY

The minimal representative of a class has the same column length distribution as its Young tableau.

PROPOSITION

Given a word w of length n , there exists an $\mathcal{O}(n^{\frac{3}{2}})$ algorithm which finds the lexicographically minimal representative equivalent to w .

COROLLARY

The minimal representative of a class has the same column length distribution as its Young tableau.

PROPOSITION

Given a word w of length n , there exists an $\mathcal{O}(n^{\frac{3}{2}})$ algorithm which finds the lexicographically minimal representative equivalent to w .

THEOREM

The set of alphabetically minimal words of the plactic congruence over an arbitrary finite alphabet is regular.

1 Cross-sections

2 Further remarks

TRANSPOSITION

Elements $x, y \in M$ are *transposed* if $x = uv$ and $y = vu$ for some $u, v \in M$.

TRANSPOSITION

Elements $x, y \in M$ are *transposed* if $x = uv$ and $y = vu$ for some $u, v \in M$.

THEOREM

Let x be an element of the plactic monoid over $\Sigma = \{a_1, \dots, a_p\}$. Define $\ell_i = |x|_{a_i}$ for $i = 1, \dots, p$. Then $(x, a_1^{\ell_1} \cdots a_p^{\ell_p}) \in T^p$.

TRANSPOSITION

Elements $x, y \in M$ are *transposed* if $x = uv$ and $y = vu$ for some $u, v \in M$.

THEOREM

Let x be an element of the plactic monoid over $\Sigma = \{a_1, \dots, a_p\}$. Define $\ell_i = |x|_{a_i}$ for $i = 1, \dots, p$. Then $(x, a_1^{\ell_1} \cdots a_p^{\ell_p}) \in T^p$.

Take the word uvw with v increasing and uw not empty and assume that

$$\max\{a \mid v \in \Sigma^* a \Sigma^*\} < b = \min\{a \mid uw \in \Sigma^* a \Sigma^*\}.$$

TRANSPOSITION

Elements $x, y \in M$ are *transposed* if $x = uv$ and $y = vu$ for some $u, v \in M$.

THEOREM

Let x be an element of the plactic monoid over $\Sigma = \{a_1, \dots, a_p\}$. Define $\ell_i = |x|_{a_i}$ for $i = 1, \dots, p$. Then $(x, a_1^{\ell_1} \cdots a_p^{\ell_p}) \in T^p$.

Take the word uvw with v increasing and uw not empty and assume that

$$\max\{a \mid v \in \Sigma^* a \Sigma^*\} < b = \min\{a \mid uw \in \Sigma^* a \Sigma^*\}.$$

Then by the Billiard Rules we have

$$vwu \equiv u'v'w'$$

where $v' = vb^r$ with $r = |uw|_b$.

TRANSPOSITION

Elements $x, y \in M$ are *transposed* if $x = uv$ and $y = vu$ for some $u, v \in M$.

THEOREM

Let x be an element of the plactic monoid over $\Sigma = \{a_1, \dots, a_p\}$. Define $\ell_i = |x|_{a_i}$ for $i = 1, \dots, p$. Then $(x, a_1^{\ell_1} \cdots a_p^{\ell_p}) \in T^p$.

CONJUGACY

Elements $x, y \in M$ are *conjugates* if exists $z \in M$ such that $xz = zy$.

PROPOSITION

In the plactic monoid we always have $C = T^{2(\|A\|-1)}$.

THANK YOU!