

Suffix conjugates of certain morphic subshifts

James D. Currie, Narad Rampersad, Kalle Saari

Department of Mathematics and Statistics
University of Winnipeg

September 16, 2013

Outline

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

1 Introduction

2 Definitions

3 The suffix conjugate

4 The suffix conjugate of the Fibonacci subshift

5 The suffix conjugate of the Thue-Morse subshift

Outline

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

1 Introduction

2 Definitions

3 The suffix conjugate

4 The suffix conjugate of the Fibonacci subshift

5 The suffix conjugate of the Thue-Morse subshift

The shift orbit closure

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

- Let A be a finite alphabet.
- Let $f: A^* \rightarrow A^*$ be a morphism and

$$\mathbf{x} = f^\omega(a) = \lim_{n \rightarrow \infty} f^n(a) \quad (a \in A).$$

- Let \mathcal{X}_f be the **shift orbit closure** of \mathbf{x} . That is,

$$\mathcal{X}_f = \{\mathbf{y} \in A^{\mathbb{N}} : \text{Fact}(\mathbf{y}) \subseteq \text{Fact}(\mathbf{x})\}$$

A representation of points in \mathcal{X}_f

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Lemma

If $\mathbf{y}_0 \in \mathcal{X}_f$, then there exist $a_0 \in A$ and $\mathbf{y}_1 \in A^{\mathbb{N}}$ such that

$$\mathbf{y}_0 = s_0 f(\mathbf{y}_1),$$

where s_0 is a nonempty suffix of $f(a_0)$ and $a_0 \mathbf{y}_1 \in \mathcal{X}_f$.

Therefore,

$$\mathbf{y}_0 = s_0 f(s_1) f^2(s_2) \cdots f^n(s_n) \cdots$$

So, every $\mathbf{y} \in \mathcal{X}_f$ has a representation as a sequence $s_0, s_1, s_2 \dots$

A representation of points in \mathcal{X}_f

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Lemma

If $\mathbf{y}_0 \in \mathcal{X}_f$, then there exist $a_0 \in A$ and $\mathbf{y}_1 \in A^{\mathbb{N}}$ such that

$$\mathbf{y}_0 = s_0 f(\mathbf{y}_1),$$

where s_0 is a nonempty suffix of $f(a_0)$ and $a_0 \mathbf{y}_1 \in \mathcal{X}_f$.

Therefore,

$$\mathbf{y}_0 = s_0 f(s_1) f^2(s_2) \cdots f^n(s_n) \cdots$$

So, every $\mathbf{y} \in \mathcal{X}_f$ has a representation as a sequence $s_0, s_1, s_2 \dots$

Problem

Which sequences s_0, s_1, s_2, \dots actually represent points in \mathcal{X}_f ?

Example

- Let $f = \varphi$ be the Fibonacci morphism $0 \mapsto 01, 1 \mapsto 0$ and \mathbf{f} the **Fibonacci word**

$$\mathbf{f} = \lim_{n \rightarrow \infty} \varphi^n(0) = 0100101001001 \dots$$

- Then

$$\begin{aligned} \mathbf{f} &= 01.0.01.010.01001.01001010. \dots \\ &= 01.\varphi^1(1).\varphi^2(1).\varphi^3(1).\dots.\varphi^n(1).\dots \end{aligned}$$

so $01, 1, 1, 1, 1 \dots$ is a representation of \mathbf{f} .

- But no sequence of suffixes that starts with $0, 1, 1, \dots$ can represent a point in \mathcal{X}_φ because

$$0.\varphi(1).\varphi^2(1).\dots = 0.0.01.\dots$$

Suffix conjugate of the Fibonacci subshift

Suffix
conjugates

CRS


Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift



Outline

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

1 Introduction

2 Definitions

3 The suffix conjugate

4 The suffix conjugate of the Fibonacci subshift

5 The suffix conjugate of the Thue-Morse subshift

The subshift (\mathcal{X}, T)

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

- Let $f : A^* \rightarrow A^*$ be a morphism.
- A fixed point $f^\omega(\alpha) = \lim_{n \rightarrow \infty} f^n(\alpha)$ for some $\alpha \in A$.
- Let \mathcal{X} be the **shift orbit closure** of $f^\omega(\alpha)$.
- The **shift map** $T : A^{\mathbb{N}} \rightarrow A^{\mathbb{N}}$ is

$$T(ax) = x \quad (a \in A)$$

- Since T is continuous and $T(\mathcal{X}) \subset \mathcal{X}$, we get a topological dynamical system (\mathcal{X}, T) called a **subshift**.

The mapping π

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

- Let $\text{Suff}(f)$ be the set of nonempty suffixes of the f -images of letters:

$$\text{Suff}(f) = \{s \in A^+ : f(a) = ps \text{ for some } a \in A\}.$$

- Denote $S = \{0, 1, \dots, |\text{Suff}(f)| - 1\}$ and let $c: S \rightarrow \text{Suff}(f)$ be a bijection.
- If $\mathbf{s} = s_0 s_1 s_2 \cdots s_n \cdots \in S^{\mathbb{N}}$ with $s_i \in S$, then we write

$$\pi(\mathbf{s}) = c(s_0)f(c(s_1))f^2(c(s_2)) \cdots f^n(c(s_n)) \cdots$$

- Thus π is a mapping $S^{\mathbb{N}} \rightarrow A^{\mathbb{N}}$.
- Define the **suffix representation** of \mathcal{X} :

$$\mathcal{S} = \{\mathbf{s} \in S^{\mathbb{N}} \mid \pi(\mathbf{s}) \in \mathcal{X}\}.$$

Example

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

- Let $f = \varphi$ with $\varphi(0) = 01$ and $\varphi(1) = 0$.
- Let $f^\omega(\alpha) = \mathbf{f}$ be the Fibonacci word.
- Then $\text{Suff}(f) = \{0, 1, 01\}$ and $S = \{0, 1, 2\}$.
- Let $c: S \rightarrow \text{Suff}(f)$ with $c(0) = 0$, $c(1) = 1$, and $c(2) = 01$.
- If $\mathbf{s} \in S^{\mathbb{N}}$ with $\mathbf{s} = 021\dots$, then

$$\begin{aligned}\pi(\mathbf{s}) &= c(0).\varphi(c(2)).\varphi^2(c(1))\dots \\ &= 0.\varphi(01).\varphi^2(1)\dots \\ &= 0.010.01\dots\end{aligned}$$

Outline

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

1 Introduction

2 Definitions

3 The suffix conjugate

4 The suffix conjugate of the Fibonacci subshift

5 The suffix conjugate of the Thue-Morse subshift

Circular morphisms

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Definition

A morphism $f: A^* \rightarrow A^*$ is **circular on** $U \subseteq A^*$ if f is injective on U and every sufficiently long $w \in A^*$ satisfies:

- 1 There exists a pair $(w_1, w_2) \in A^* \times A^*$ such that

$$w = w_1 w_2$$

- 2 For all $v_1, v_2 \in A^*$,

$$v_1 w v_2 \in f(U) \implies v_1 w_1 \in f(U) \quad \text{and} \quad w_2 v_2 \in f(U).$$

Example

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

- The Fibonacci morphism $\varphi: \{0, 1\}^* \rightarrow \{0, 1\}^*$ with

$$\varphi(0) = 01 \quad \text{and} \quad \varphi(1) = 0$$

is circular on $\{0, 1\}^*$.

- The Thue-Morse morphism $\mu: \{0, 1\}^* \rightarrow \{0, 1\}^*$ with

$$\mu(0) = 01 \quad \text{and} \quad \mu(1) = 10$$

is **not** circular on $\{0, 1\}^*$ but it is circular on $\text{Fact}(\mathbf{t})$,
where \mathbf{t} is the Thue-Morse word.

Mossé's Theorem

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Theorem (Mossé, 1991)

*If f is a **primitive** morphism with an **aperiodic** fixed point \mathbf{x} , then f is **circular** on $\text{Fact}(\mathbf{x})$.*

The morphism class \mathcal{N}

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Definition

Let $f: A^* \rightarrow A^*$ be a morphism and let $\mathbf{x} = f^\omega(a)$ ($a \in A$) be aperiodic. Then we say that f is in class \mathcal{N} if

- 1 f is circular on $\text{Fact}(\mathbf{x})$.
- 2 the set $f(A)$ is a suffix code.
- 3 for every $a \in A$, we have $|f^n(a)| \rightarrow \infty$ as $n \rightarrow \infty$.

Remark

If f is primitive, \mathbf{x} aperiodic, and $f(A)$ a suffix code, then $f \in \mathcal{N}$ by Mossé's Theorem.

Example

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Example

- The Fibonacci morphism $\varphi(0) = 01, \varphi(1) = 0$ is primitive.
- $\varphi(\{0, 1\}) = \{01, 1\}$ is a suffix code.
- $\mathbf{f} = \varphi^\omega(0)$ is aperiodic.
- Thus $\varphi \in \mathcal{N}$.

Example

- The Thue-Morse morphism $\mu(0) = 01, \mu(1) = 10$ is primitive.
- $\mu(\{0, 1\}) = \{01, 10\}$ is a suffix code.
- $\mu^\omega(0)$ is aperiodic.
- Thus $\mu \in \mathcal{N}$.

Motivating class \mathcal{N}

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Recall the suffix representation

$$\mathcal{S} = \{ \mathbf{s} \in S^{\mathbb{N}} \mid \pi(\mathbf{s}) \in \mathcal{X} \}.$$

and the shift map $T(a\mathbf{x}) = \mathbf{x} \quad (a \in A)$.

Lemma

If $f \in \mathcal{N}$, then $T(\mathcal{S}) \subseteq \mathcal{S}$. Thus (\mathcal{S}, T) is a subshift.

Lemma

If $f \in \mathcal{N}$, then $\pi: \mathcal{S} \rightarrow \mathcal{X}$ is injective.

The mappings λ and G

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Definition

Let $\lambda: S^* \rightarrow S^*$ be the morphism given by

$$\lambda(s) = \begin{cases} c^{-1}(f(c(s))) & \text{if } c(s) \in A; \\ c^{-1}(f(a))c^{-1}(u) & \text{if } c(s) = au \text{ with } a \in A \text{ and } u \in A^+ \end{cases}$$

Write

$$S_1 = \{s \in S : c(s) \in A\}.$$

Definition

Let $G: S^{\mathbb{N}} \rightarrow S^{\mathbb{N}}$ be defined by

$$G(s) = \begin{cases} \lambda(ps)t & \text{if } s = pst \text{ with } p \in S_1^* \text{ and } s \in S \setminus S_1 \\ \lambda(s) & \text{if } s \in S_1^{\mathbb{N}}. \end{cases}$$



The Suffix Conjugate

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate


The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Let $H: S \rightarrow S$ be the mapping defined by $H = T \circ G$.

Theorem (Currie, Rampersad, S. 2013)

If $f \in \mathcal{N}$, then $\pi \circ H = T \circ \pi$ and $\pi: (S, H) \rightarrow (\mathcal{X}, T)$ is a conjugacy.



Definition

If $f \in \mathcal{N}$, we call (S, H) the **suffix conjugate** of (\mathcal{X}, T)

Outline

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

1 Introduction

2 Definitions

3 The suffix conjugate

4 The suffix conjugate of the Fibonacci subshift

5 The suffix conjugate of the Thue-Morse subshift

The mapping H_φ

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Lemma

The mapping $H_\varphi: \mathcal{S}_\varphi \rightarrow \mathcal{S}_\varphi$ of the suffix conjugate $(\mathcal{S}_\varphi, H_\varphi)$ of the Fibonacci subshift (\mathcal{X}_φ, T) is given by

$$H_\varphi(\mathbf{s}) = \begin{cases} 1\mathbf{z} & \text{if } \mathbf{s} = 2\mathbf{z}; \\ \lambda(x2)\mathbf{z} & \text{if } \mathbf{s} = ax2\mathbf{z} \text{ with } a \in \{0, 1\}, x \in \{0, 1\}^*; \\ \lambda(\mathbf{z}) & \text{if } \mathbf{s} = a\mathbf{z} \text{ with } a \in \{0, 1\} \text{ and } \mathbf{z} \in \{0, 1\}^{\mathbb{N}}, \end{cases}$$

where λ is the morphism

$$\lambda(1) = 0, \quad \lambda(0) = 2, \quad \text{and} \quad \lambda(2) = 21.$$

The language of the suffix conjugate

Suffix
conjugates

CRS

Introduction

Definitions


The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Theorem (Currie, Rampersad, S. 2013)

The language $\mathcal{L}(\mathcal{S}_\varphi)$ of the suffix conjugate $(\mathcal{S}_\varphi, H_\varphi)$ of the Fibonacci subshift (\mathcal{X}_φ, T) is regular. An infinite word $\mathbf{s} \in S^{\mathbb{N}}$ is in \mathcal{S}_φ if and only if it is the label of an infinite walk on the graph below.



Example

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Let $\mathbf{f} \in \mathcal{X}_\varphi$ be the Fibonacci word. Then

$$\blacksquare \quad \mathbf{s} = 2121^\omega \xrightarrow{\pi} 010\mathbf{f}$$

Example

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Let $\mathbf{f} \in \mathcal{X}_\varphi$ be the Fibonacci word. Then

$$\blacksquare \quad \mathbf{s} = 2121^\omega \quad \xrightarrow{\pi} \quad 010\mathbf{f}$$

$$\blacksquare \quad H_\varphi(\mathbf{s}) = 1121^\omega \quad \xrightarrow{\pi} \quad 10\mathbf{f}$$

Example

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Let $\mathbf{f} \in \mathcal{X}_\varphi$ be the Fibonacci word. Then

- $\mathbf{s} = 2121^\omega \xrightarrow{\pi} 010\mathbf{f}$
- $H_\varphi(\mathbf{s}) = 1121^\omega \xrightarrow{\pi} 10\mathbf{f}$
- $H_\varphi^2(\mathbf{s}) = 021^\omega \xrightarrow{\pi} 0\mathbf{f}$

Example

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Let $\mathbf{f} \in \mathcal{X}_\varphi$ be the Fibonacci word. Then

$$\blacksquare \quad \mathbf{s} = 2121^\omega \quad \xrightarrow{\pi} \quad 010\mathbf{f}$$

$$\blacksquare \quad H_\varphi(\mathbf{s}) = 1121^\omega \quad \xrightarrow{\pi} \quad 10\mathbf{f}$$

$$\blacksquare \quad H_\varphi^2(\mathbf{s}) = 021^\omega \quad \xrightarrow{\pi} \quad 0\mathbf{f}$$

$$\blacksquare \quad H_\varphi^3(\mathbf{s}) = 21^\omega \quad \xrightarrow{\pi} \quad \mathbf{f}$$

Example

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Let $\mathbf{f} \in \mathcal{X}_\varphi$ be the Fibonacci word. Then

- $\mathbf{s} = 2121^\omega \xrightarrow{\pi} 010\mathbf{f}$
- $H_\varphi(\mathbf{s}) = 1121^\omega \xrightarrow{\pi} 10\mathbf{f}$
- $H_\varphi^2(\mathbf{s}) = 021^\omega \xrightarrow{\pi} 0\mathbf{f}$
- $H_\varphi^3(\mathbf{s}) = 21^\omega \xrightarrow{\pi} \mathbf{f}$
- $H_\varphi^4(\mathbf{s}) = 1^\omega \xrightarrow{\pi} T\mathbf{f}$

Example

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Let $\mathbf{f} \in \mathcal{X}_\varphi$ be the Fibonacci word. Then

- $\mathbf{s} = 2121^\omega \xrightarrow{\pi} 010\mathbf{f}$
- $H_\varphi(\mathbf{s}) = 1121^\omega \xrightarrow{\pi} 10\mathbf{f}$
- $H_\varphi^2(\mathbf{s}) = 021^\omega \xrightarrow{\pi} 0\mathbf{f}$
- $H_\varphi^3(\mathbf{s}) = 21^\omega \xrightarrow{\pi} \mathbf{f}$
- $H_\varphi^4(\mathbf{s}) = 1^\omega \xrightarrow{\pi} T\mathbf{f}$
- $H_\varphi^5(\mathbf{s}) = 0^\omega \xrightarrow{\pi} T^2\mathbf{f}$

Example

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Let $\mathbf{f} \in \mathcal{X}_\varphi$ be the Fibonacci word. Then

- $\mathbf{s} = 2121^\omega \xrightarrow{\pi} 010\mathbf{f}$
- $H_\varphi(\mathbf{s}) = 1121^\omega \xrightarrow{\pi} 10\mathbf{f}$
- $H_\varphi^2(\mathbf{s}) = 021^\omega \xrightarrow{\pi} 0\mathbf{f}$
- $H_\varphi^3(\mathbf{s}) = 21^\omega \xrightarrow{\pi} \mathbf{f}$
- $H_\varphi^4(\mathbf{s}) = 1^\omega \xrightarrow{\pi} T\mathbf{f}$
- $H_\varphi^5(\mathbf{s}) = 0^\omega \xrightarrow{\pi} T^2\mathbf{f}$
- $H_\varphi^6(\mathbf{s}) = 2^\omega \xrightarrow{\pi} T^3\mathbf{f}$

Example

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Let $\mathbf{f} \in \mathcal{X}_\varphi$ be the Fibonacci word. Then

- $\mathbf{s} = 2121^\omega \xrightarrow{\pi} 010\mathbf{f}$
- $H_\varphi(\mathbf{s}) = 1121^\omega \xrightarrow{\pi} 10\mathbf{f}$
- $H_\varphi^2(\mathbf{s}) = 021^\omega \xrightarrow{\pi} 0\mathbf{f}$
- $H_\varphi^3(\mathbf{s}) = 21^\omega \xrightarrow{\pi} \mathbf{f}$
- $H_\varphi^4(\mathbf{s}) = 1^\omega \xrightarrow{\pi} T\mathbf{f}$
- $H_\varphi^5(\mathbf{s}) = 0^\omega \xrightarrow{\pi} T^2\mathbf{f}$
- $H_\varphi^6(\mathbf{s}) = 2^\omega \xrightarrow{\pi} T^3\mathbf{f}$
- $H_\varphi^7(\mathbf{s}) = 12^\omega \xrightarrow{\pi} T^4\mathbf{f}$

Points in the orbit of the Fibonacci word

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

- An infinite word $\mathbf{y} \in A^{\mathbb{N}}$ is a **tail** (or **suffix**) of $\mathbf{x} \in A^{\mathbb{N}}$ if $T^n \mathbf{x} = \mathbf{y}$ for some $n \geq 0$.
- The **orbit** of $\mathbf{x} \in A^{\mathbb{N}}$ is the set

$$\{\mathbf{y} \in A^{\mathbb{N}} : \mathbf{x} \text{ is a tail of } \mathbf{y} \text{ or vice versa}\}$$

Theorem (Currie, Rampersad, S. 2013)

$T^2 \mathbf{f}$ splits the orbit of the Fibonacci word \mathbf{f} in the following way: Let $\mathbf{s} \in \mathcal{S}_\varphi$.

- *$T^2 \mathbf{f}$ is a proper tail of $\pi(\mathbf{s}) \iff 1^\omega$ is a tail of \mathbf{s} .*
- *$\pi(\mathbf{s})$ is a proper tail of $T^2 \mathbf{f} \iff 2^\omega$ is a tail of \mathbf{s} .*

Incidentally, $\pi(0^\omega) = T^2 \mathbf{f}$.

Outline

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

1 Introduction

2 Definitions

3 The suffix conjugate

4 The suffix conjugate of the Fibonacci subshift

5 The suffix conjugate of the Thue-Morse subshift

The mapping H_μ

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Lemma

The mapping $H_\mu: S_\mu \rightarrow S_\mu$ of the suffix conjugate (S_μ, H_μ) of the Thue-Morse subshift (\mathcal{X}_μ, T) is given by

$$H_\mu(\mathbf{s}) = \begin{cases} 1\mathbf{z} & \text{if } \mathbf{s} = 2\mathbf{z}; \\ 0\mathbf{z} & \text{if } \mathbf{s} = 3\mathbf{z}; \\ \lambda(\mathbf{z}) & \text{if } \mathbf{s} = a\mathbf{z} \text{ with } a \in \{0, 1\} \text{ and } \mathbf{z} \in \{0, 1\}^{\mathbb{N}}; \\ \lambda(x2)\mathbf{z} & \text{if } \mathbf{s} = ax2\mathbf{z} \text{ with } a \in \{0, 1\}, x \in \{0, 1\}^*; \\ \lambda(x3)\mathbf{z} & \text{if } \mathbf{s} = ax3\mathbf{z} \text{ with } a \in \{0, 1\}, x \in \{0, 1\}^*, \end{cases}$$


where λ is the morphism given by

$$\lambda(0) = 2, \quad \lambda(1) = 3, \quad \lambda(2) = 21, \quad \text{and} \quad \lambda(3) = 30.$$

The language of \mathcal{S}_μ

Theorem (Currie, Rampersad, S. 2013)

The language $\mathcal{L}(\mathcal{S}_\mu)$ of the suffix conjugate (\mathcal{S}_μ, H_μ) of the Thue-Morse subshift (\mathcal{X}_μ, T) is regular. An infinite word $\mathbf{s} \in S^{\mathbb{N}}$ is in \mathcal{S}_μ if and only if it is the label of an infinite walk on the graph below.



Points in the orbit of the Thue-Morse word

Suffix
conjugates

CRS

Introduction

Definitions

The suffix
conjugate

The suffix
conjugate of
the Fibonacci
subshift

The suffix
conjugate of
the
Thue-Morse
subshift

Theorem (Currie, Rampersad, S. 2013)

Let $\mathbf{s} \in \mathcal{S}_\mu$. Then

- (i) $\pi(\mathbf{s})$ is a proper tail of $T^2\mathbf{t}$ if and only if 3^ω is a tail of \mathbf{s} .*
- (ii) $T\mathbf{t}$ is a proper tail of $\pi(\mathbf{s})$ if and only if 1^ω is a tail of \mathbf{s} .*

Thank you!