

# Similarity Relations and Repetition-Freeness

Tomi Kärki

Department of Teacher Education and  
Department of Mathematics and Statistics,  
University of Turku, Finland

9th International Conference on WORDS 2013  
Turku, September 16-20, 2013

- 1 Background
- 2 Similarity relation
- 3 Relational repetition-freeness
- 4 Square-freeness and overlap-freeness
- 5 Cube-freeness and  $3^+$ -freeness

- seminal papers of Thue (1906, 1912)

- seminal papers of Thue (1906, 1912)
- abelian repetitions: Erdős (1961), Dekking (1979), Keränen (1992)

- seminal papers of Thue (1906, 1912)
- abelian repetitions: Erdős (1961), Dekking (1979), Keränen (1992)
- $c$ -approximate and  $\alpha$ -similar squares: Krieger, Ochermann, Rampersad, Shallit (2007)

- seminal papers of Thue (1906, 1912)
- abelian repetitions: Erdős (1961), Dekking (1979), Keränen (1992)
- $c$ -approximate and  $\alpha$ -similar squares: Krieger, Ochermann, Rampersad, Shallit (2007)
- repetitions in partial words: Manea, Mercaş, Blanchet-Sadri, ... (2007- )

- seminal papers of Thue (1906, 1912)
- abelian repetitions: Erdős (1961), Dekking (1979), Keränen (1992)
- $c$ -approximate and  $\alpha$ -similar squares: Krieger, Ochermann, Rampersad, Shallit (2007)
- repetitions in partial words: Manea, Mercaş, Blanchet-Sadri, ... (2007- )
- similarity relations: Halava, Harju, Kärki

# Similarity relation

## Definitions

- Let  $\mathcal{A}$  be an alphabet.
- Let  $R_{\mathcal{A}} \subseteq \mathcal{A} \times \mathcal{A}$  be a *reflexive* and *symmetric* relation.



# Similarity relation

## Definitions

- Let  $\mathcal{A}$  be an alphabet.
- Let  $R_{\mathcal{A}} \subseteq \mathcal{A} \times \mathcal{A}$  be a *reflexive* and *symmetric* relation.
- Two words  $u = u_1 \cdots u_n$  and  $v = v_1 \cdots v_m$ ,  $u_i, v_j \in \mathcal{A}$  are *R-similar* if  $n = m$  and  $(u_i, v_i) \in R_{\mathcal{A}}$ . Then we denote  $u R v$ .

# Similarity relation

## Definitions

- Let  $\mathcal{A}$  be an alphabet.
- Let  $R_{\mathcal{A}} \subseteq \mathcal{A} \times \mathcal{A}$  be a *reflexive* and *symmetric* relation.
- Two words  $u = u_1 \cdots u_n$  and  $v = v_1 \cdots v_m$ ,  $u_i, v_j \in \mathcal{A}$  are *R-similar* if  $n = m$  and  $(u_i, v_i) \in R_{\mathcal{A}}$ . Then we denote  $u R v$ .
- A similarity relation  $R$  can be represented by listing the  $R$ -similar letters:

$$R = \langle \{r_1, \dots, r_n\} \rangle, \quad r_i = (a_i, b_i) \in R_{\mathcal{A}}.$$

# Similarity relation

## Definitions


- Let  $\mathcal{A}$  be an alphabet.
- Let  $R_{\mathcal{A}} \subseteq \mathcal{A} \times \mathcal{A}$  be a *reflexive* and *symmetric* relation.
- Two words  $u = u_1 \cdots u_n$  and  $v = v_1 \cdots v_m$ ,  $u_i, v_j \in \mathcal{A}$  are *R-similar* if  $n = m$  and  $(u_i, v_i) \in R_{\mathcal{A}}$ . Then we denote  $u R v$ .
- A similarity relation  $R$  can be represented by listing the  $R$ -similar letters:

$$R = \langle \{r_1, \dots, r_n\} \rangle, \quad r_i = (a_i, b_i) \in R_{\mathcal{A}}.$$

- A similarity relation  $R$  can be represented by an undirected graph  $G_R = (V, E)$ , where  $V = \mathcal{A}$  and  $(a, b) \in E$  iff  $a R b$ .

# Similarity relation

## Example




## Example

- $R = \langle (a, b), (b, c), (c, d), (c, e) \rangle$

# Similarity relation

## Example




## Example

- $R = \langle (a, b), (b, c), (c, d), (c, e) \rangle$
- $abcde R acccc$

# Similarity relation

## Example




## Example

- $R = \langle (a, b), (b, c), (c, d), (c, e) \rangle$
- $abcde R acccc$

# Similarity relation

## Example




## Example

- $R = \langle (a, b), (b, c), (c, d), (c, e) \rangle$
- $abcde R a\textcolor{red}{c}ccc$

# Similarity relation

## Example




## Example

- $R = \langle (a, b), (b, c), (c, d), (c, e) \rangle$
- $ab\textcolor{red}{c}de \ R \ ac\textcolor{red}{c}cc$



# Similarity relation

## Example




## Example

- $R = \langle (a, b), (b, c), (c, d), (c, e) \rangle$
- $abc\textcolor{red}{d}e \ R \ acc\textcolor{red}{c}c$

# Similarity relation

## Example




## Example

- $R = \langle (a, b), (b, c), (c, d), (c, e) \rangle$
- $abcd\textcolor{red}{e} R a\textcolor{red}{c}cc\textcolor{red}{c}$

# Similarity relation

## Example




### Example

- $R = \langle (a, b), (b, c), (c, d), (c, e) \rangle$
- $abcde R acccc$

# Similarity relation

## Example



## Example

- $R = \langle (a, b), (b, c), (c, d), (c, e) \rangle$
- $abcde R acccc$
- $(abcde, ccccc) \notin R$

- Partial words were invented by Berstel and Boasson (1999)

# Similarity relation

## Partial words

- Partial words were invented by Berstel and Boasson (1999)
- They are words over an extended alphabet  $\mathcal{A}_\diamond = \mathcal{A} \cup \{\diamond\}$

# Similarity relation

## Partial words

- Partial words were invented by Berstel and Boasson (1999)
- They are words over an extended alphabet  $\mathcal{A}_\diamond = \mathcal{A} \cup \{\diamond\}$
- The letter  $\diamond$  is a *hole*, i.e., a 'do not know'-symbol which may correspond to any letter of the alphabet  $\mathcal{A}$

# Similarity relation

## Partial words

- Partial words were invented by Berstel and Boasson (1999)
- They are words over an extended alphabet  $\mathcal{A}_\diamond = \mathcal{A} \cup \{\diamond\}$
- The letter  $\diamond$  is a *hole*, i.e., a 'do not know'-symbol which may correspond to any letter of the alphabet  $\mathcal{A}$
- Two partial words  $u = u_1 \cdots u_n$  and  $v = v_1 \cdots v_n$ ,  $u_i, v_i \in \mathcal{A}$  are *compatible* if  $u_i = v_i$  whenever both  $u_i$  and  $v_i$  are not holes. Then we denote  $u \uparrow v$ .



# Similarity relation

## Partial words

- Partial words were invented by Berstel and Boasson (1999)
- They are words over an extended alphabet  $\mathcal{A}_\diamond = \mathcal{A} \cup \{\diamond\}$
- The letter  $\diamond$  is a *hole*, i.e., a 'do not know'-symbol which may correspond to any letter of the alphabet  $\mathcal{A}$
- Two partial words  $u = u_1 \cdots u_n$  and  $v = v_1 \cdots v_n$ ,  $u_i, v_i \in \mathcal{A}$  are *compatible* if  $u_i = v_i$  whenever both  $u_i$  and  $v_i$  are not holes. Then we denote  $u \uparrow v$ .
- $a \diamond c b \diamond \uparrow \diamond b c b \diamond$

# Similarity relation

## Partial words

- Partial words were invented by Berstel and Boasson (1999)
- They are words over an extended alphabet  $\mathcal{A}_\diamond = \mathcal{A} \cup \{\diamond\}$
- The letter  $\diamond$  is a *hole*, i.e., a 'do not know'-symbol which may correspond to any letter of the alphabet  $\mathcal{A}$
- Two partial words  $u = u_1 \cdots u_n$  and  $v = v_1 \cdots v_n$ ,  $u_i, v_i \in \mathcal{A}$  are *compatible* if  $u_i = v_i$  whenever both  $u_i$  and  $v_i$  are not holes. Then we denote  $u \uparrow v$ .
- $a \diamond cb \diamond \uparrow \diamond bcb \diamond$

# Similarity relation

## Partial words

- Partial words were invented by Berstel and Boasson (1999)
- They are words over an extended alphabet  $\mathcal{A}_\diamond = \mathcal{A} \cup \{\diamond\}$
- The letter  $\diamond$  is a *hole*, i.e., a 'do not know'-symbol which may correspond to any letter of the alphabet  $\mathcal{A}$
- Two partial words  $u = u_1 \cdots u_n$  and  $v = v_1 \cdots v_n$ ,  $u_i, v_i \in \mathcal{A}$  are *compatible* if  $u_i = v_i$  whenever both  $u_i$  and  $v_i$  are not holes. Then we denote  $u \uparrow v$ .
- $a \diamond cb \diamond \uparrow \diamond bcb \diamond$

# Similarity relation

## Partial words

- Partial words were invented by Berstel and Boasson (1999)
- They are words over an extended alphabet  $\mathcal{A}_\diamond = \mathcal{A} \cup \{\diamond\}$
- The letter  $\diamond$  is a *hole*, i.e., a 'do not know'-symbol which may correspond to any letter of the alphabet  $\mathcal{A}$
- Two partial words  $u = u_1 \cdots u_n$  and  $v = v_1 \cdots v_n$ ,  $u_i, v_i \in \mathcal{A}$  are *compatible* if  $u_i = v_i$  whenever both  $u_i$  and  $v_i$  are not holes. Then we denote  $u \uparrow v$ .
- $a \diamond c b \diamond \uparrow \diamond b c b \diamond$

# Similarity relation

## Partial words

- Partial words were invented by Berstel and Boasson (1999)
- They are words over an extended alphabet  $\mathcal{A}_\diamond = \mathcal{A} \cup \{\diamond\}$
- The letter  $\diamond$  is a *hole*, i.e., a 'do not know'-symbol which may correspond to any letter of the alphabet  $\mathcal{A}$
- Two partial words  $u = u_1 \cdots u_n$  and  $v = v_1 \cdots v_n$ ,  $u_i, v_i \in \mathcal{A}$  are *compatible* if  $u_i = v_i$  whenever both  $u_i$  and  $v_i$  are not holes. Then we denote  $u \uparrow v$ .
- $a \diamond c \textcolor{red}{b} \diamond \uparrow \diamond b c \textcolor{red}{b} \diamond$

# Similarity relation

## Partial words

- Partial words were invented by Berstel and Boasson (1999)
- They are words over an extended alphabet  $\mathcal{A}_\diamond = \mathcal{A} \cup \{\diamond\}$
- The letter  $\diamond$  is a *hole*, i.e., a 'do not know'-symbol which may correspond to any letter of the alphabet  $\mathcal{A}$
- Two partial words  $u = u_1 \cdots u_n$  and  $v = v_1 \cdots v_n$ ,  $u_i, v_i \in \mathcal{A}$  are *compatible* if  $u_i = v_i$  whenever both  $u_i$  and  $v_i$  are not holes. Then we denote  $u \uparrow v$ .
- $a \diamond cb \uparrow \diamond bcb \diamond$

# Similarity relation

## Partial words

- Partial words were invented by Berstel and Boasson (1999)
- They are words over an extended alphabet  $\mathcal{A}_\diamond = \mathcal{A} \cup \{\diamond\}$
- The letter  $\diamond$  is a *hole*, i.e., a 'do not know'-symbol which may correspond to any letter of the alphabet  $\mathcal{A}$
- Two partial words  $u = u_1 \cdots u_n$  and  $v = v_1 \cdots v_n$ ,  $u_i, v_i \in \mathcal{A}$  are *compatible* if  $u_i = v_i$  whenever both  $u_i$  and  $v_i$  are not holes. Then we denote  $u \uparrow v$ .
- $a \diamond cb \diamond \uparrow \diamond bcb \diamond$ ,

# Similarity relation

## Partial words

- Partial words were invented by Berstel and Boasson (1999)
- They are words over an extended alphabet  $\mathcal{A}_\diamond = \mathcal{A} \cup \{\diamond\}$
- The letter  $\diamond$  is a *hole*, i.e., a 'do not know'-symbol which may correspond to any letter of the alphabet  $\mathcal{A}$
- Two partial words  $u = u_1 \cdots u_n$  and  $v = v_1 \cdots v_n$ ,  $u_i, v_i \in \mathcal{A}$  are *compatible* if  $u_i = v_i$  whenever both  $u_i$  and  $v_i$  are not holes. Then we denote  $u \uparrow v$ .
- $a \diamond cb \diamond \uparrow \diamond bcb \diamond$ ,  $a \diamond cb \diamond \not\uparrow b \diamond cba$



# Similarity relation

## Partial words

- Partial words were invented by Berstel and Boasson (1999)
- They are words over an extended alphabet  $\mathcal{A}_\diamond = \mathcal{A} \cup \{\diamond\}$
- The letter  $\diamond$  is a *hole*, i.e., a 'do not know'-symbol which may correspond to any letter of the alphabet  $\mathcal{A}$
- Two partial words  $u = u_1 \cdots u_n$  and  $v = v_1 \cdots v_n$ ,  $u_i, v_i \in \mathcal{A}$  are *compatible* if  $u_i = v_i$  whenever both  $u_i$  and  $v_i$  are not holes. Then we denote  $u \uparrow v$ .
- $a \diamond cb \diamond \uparrow \diamond bcb \diamond$ ,  $a \diamond cb \diamond \not\uparrow b \diamond cba$

# Similarity relation

## Partial words

- Partial words were invented by Berstel and Boasson (1999)
- They are words over an extended alphabet  $\mathcal{A}_\diamond = \mathcal{A} \cup \{\diamond\}$
- The letter  $\diamond$  is a *hole*, i.e., a 'do not know'-symbol which may correspond to any letter of the alphabet  $\mathcal{A}$
- Two partial words  $u = u_1 \cdots u_n$  and  $v = v_1 \cdots v_n$ ,  $u_i, v_i \in \mathcal{A}$  are *compatible* if  $u_i = v_i$  whenever both  $u_i$  and  $v_i$  are not holes. Then we denote  $u \uparrow v$ .
- $a \diamond cb \diamond \uparrow \diamond bcb \diamond$ ,  $a \diamond cb \diamond \not\uparrow b \diamond cba$
- Compatibility relation of partial words corresponds to the similarity relation  $R_\uparrow = \langle \{(\diamond, a) \mid a \in \mathcal{A}\} \rangle$ .

# Similarity relation

## Chain relations and cyclic relations

- Let  $\mathcal{A}_n = \{0, 1, \dots, n-1\}$ .
- *Chain relation*  $\overline{R}_n$  is a similarity relation on  $\mathcal{A}_n^*$  defined by  $\overline{R}_n = \langle \{(i, i+1) \mid i = 0, 1, \dots, n-2\} \rangle$
- *Cyclic relation*  $\mathring{R}_n$  is a similarity relation on  $\mathcal{A}_n^*$  defined by  $\mathring{R}_n = \langle \{(i, i+1) \mid i = 0, 1, \dots, n-2\} \cup \{(n-1, 0)\} \rangle$ .

# Similarity relation


## Chain relations and cyclic relations

- Let  $\mathcal{A}_n = \{0, 1, \dots, n-1\}$ .
- *Chain relation*  $\overline{R}_n$  is a similarity relation on  $\mathcal{A}_n^*$  defined by  $\overline{R}_n = \langle \{(i, i+1) \mid i = 0, 1, \dots, n-2\} \rangle$
- *Cyclic relation*  $\dot{R}_n$  is a similarity relation on  $\mathcal{A}_n^*$  defined by  $\dot{R}_n = \langle \{(i, i+1) \mid i = 0, 1, \dots, n-2\} \cup \{(n-1, 0)\} \rangle$ .
- 543210  $\dot{R}_6$  432105
- $(543210, 432105) \notin \overline{R}_6$


$G_{\overline{R}_6} :$

0 — 1 — 2 — 3 — 4 — 5

$G_{\dot{R}_6} :$



$G_{R_{\uparrow}} :$



# Relational repetition-freeness

Global and local  $R$ -repetitions

- Let  $\text{pref}_\ell(w)$  denote the prefix of length  $\ell$  of a word  $w$ .

# Relational repetition-freeness

## Global and local $R$ -repetitions

- Let  $\text{pref}_\ell(w)$  denote the prefix of length  $\ell$  of a word  $w$ .
- A word  $w \in \mathcal{A}^+$  is a *global  $R$ -repetition of order  $k$*  if  $w = u_1 \cdots u_m u'$  where the words  $u_i$  are pairwise  $R$ -similar,  $u' R \text{pref}_{|u'|}(u_i)$  for every  $i = 1, 2, \dots, m$  and  $k = |w|/|u_i|$ .

# Relational repetition-freeness

## Global and local $R$ -repetitions

- Let  $\text{pref}_\ell(w)$  denote the prefix of length  $\ell$  of a word  $w$ .
- A word  $w \in \mathcal{A}^+$  is a *global  $R$ -repetition of order  $k$*  if  $w = u_1 \cdots u_m u'$  where the words  $u_i$  are pairwise  $R$ -similar,  $u' R \text{pref}_{|u'|}(\text{pref}_{|u'|}(u_i))$  for every  $i = 1, 2, \dots, m$  and  $k = |w|/|u_i|$ .
- A word  $w \in \mathcal{A}^+$  is a *local  $R$ -repetition of order  $k$*  if  $w = u_1 \cdots u_m u'$  where  $u_i R u_{i+1}$  for  $i = 1, 2, \dots, m-1$ ,  $u' R \text{pref}_{|u'|}(u_m)$  and  $k = |w|/|u_i|$ .

# Relational repetition-freeness

## Global and local $R$ -repetitions

- Let  $\text{pref}_\ell(w)$  denote the prefix of length  $\ell$  of a word  $w$ .
- A word  $w \in \mathcal{A}^+$  is a *global  $R$ -repetition of order  $k$*  if  $w = u_1 \cdots u_m u'$  where the words  $u_i$  are pairwise  $R$ -similar,  $u' R \text{pref}_{|u'|}(\text{pref}_{|u'|}(u_i))$  for every  $i = 1, 2, \dots, m$  and  $k = |w|/|u_i|$ .
- A word  $w \in \mathcal{A}^+$  is a *local  $R$ -repetition of order  $k$*  if  $w = u_1 \cdots u_m u'$  where  $u_i R u_{i+1}$  for  $i = 1, 2, \dots, m-1$ ,  $u' R \text{pref}_{|u'|}(u_m)$  and  $k = |w|/|u_i|$ .

### Example

- $\overline{R}_3 = \langle (0, 1), (1, 2) \rangle$
- $u = \text{0021022}$  is a local  $\overline{R}_3$ -repetition of order  $\frac{7}{3}$



# Relational repetition-freeness

## Global and local $R$ -repetitions

- Let  $\text{pref}_\ell(w)$  denote the prefix of length  $\ell$  of a word  $w$ .
- A word  $w \in \mathcal{A}^+$  is a *global  $R$ -repetition of order  $k$*  if  $w = u_1 \cdots u_m u'$  where the words  $u_i$  are pairwise  $R$ -similar,  $u' R \text{pref}_{|u'|}(u_i)$  for every  $i = 1, 2, \dots, m$  and  $k = |w|/|u_i|$ .
- A word  $w \in \mathcal{A}^+$  is a *local  $R$ -repetition of order  $k$*  if  $w = u_1 \cdots u_m u'$  where  $u_i R u_{i+1}$  for  $i = 1, 2, \dots, m-1$ ,  $u' R \text{pref}_{|u'|}(u_m)$  and  $k = |w|/|u_i|$ .

### Example

- $\overline{R}_3 = \langle (0, 1), (1, 2) \rangle$
- $u = \text{0021022}$  is a local  $\overline{R}_3$ -repetition of order  $\frac{7}{3}$
- $u = \text{0021022}$  is not a global  $\overline{R}_3$ -repetition of order  $\frac{7}{3}$

- A word  $w \in \mathcal{A}^+$  is *globally* (resp. *locally*)  $(R, k)$ -free if it does not contain any global (resp. local)  $R$ -repetitions of order  $k$ .

- A word  $w \in \mathcal{A}^+$  is *globally* (resp. *locally*)  $(R, k)$ -free if it does not contain any global (resp. local)  $R$ -repetitions of order  $k$ .
- A word  $w \in \mathcal{A}^+$  is *globally* (resp. *locally*)  $(R, k^+)$ -free if it does not contain any global (resp. local)  $R$ -repetitions of order  $\ell > k$ .

- A word  $w \in \mathcal{A}^+$  is *globally* (resp. *locally*)  $(R, k)$ -free if it does not contain any global (resp. local)  $R$ -repetitions of order  $k$ .
- A word  $w \in \mathcal{A}^+$  is *globally* (resp. *locally*)  $(R, k^+)$ -free if it does not contain any global (resp. local)  $R$ -repetitions of order  $\ell > k$ .
- An  $R$ -repetition of order 2 is called an  *$R$ -square*.

- A word  $w \in \mathcal{A}^+$  is *globally* (resp. *locally*)  $(R, k)$ -free if it does not contain any global (resp. local)  $R$ -repetitions of order  $k$ .
- A word  $w \in \mathcal{A}^+$  is *globally* (resp. *locally*)  $(R, k^+)$ -free if it does not contain any global (resp. local)  $R$ -repetitions of order  $\ell > k$ .
- An  $R$ -repetition of order 2 is called an  *$R$ -square*.
- An  $R$ -repetition of order 3 is called an  *$R$ -cube*.

- A word  $w \in \mathcal{A}^+$  is *globally* (resp. *locally*)  $(R, k)$ -free if it does not contain any global (resp. local)  $R$ -repetitions of order  $k$ .
- A word  $w \in \mathcal{A}^+$  is *globally* (resp. *locally*)  $(R, k^+)$ -free if it does not contain any global (resp. local)  $R$ -repetitions of order  $\ell > k$ .
- An  $R$ -repetition of order 2 is called an  *$R$ -square*.
- An  $R$ -repetition of order 3 is called an  *$R$ -cube*.
- A *local  $R$ -overlap* is a word of the form  $uu'vv'w$ , where  $u R v$ ,  $u' R v'$  and  $v R w$ . In this case, there are two  $R$ -similar overlapping words  $uu'v$  and  $vv'w$ .

- A word  $w \in \mathcal{A}^+$  is *globally* (resp. *locally*)  $(R, k)$ -free if it does not contain any global (resp. local)  $R$ -repetitions of order  $k$ .
- A word  $w \in \mathcal{A}^+$  is *globally* (resp. *locally*)  $(R, k^+)$ -free if it does not contain any global (resp. local)  $R$ -repetitions of order  $\ell > k$ .
- An  $R$ -repetition of order 2 is called an  *$R$ -square*.
- An  $R$ -repetition of order 3 is called an  *$R$ -cube*.
- A *local  $R$ -overlap* is a word of the form  $uu'vv'w$ , where  $u R v$ ,  $u' R v'$  and  $v R w$ . In this case, there are two  $R$ -similar overlapping words  $uu'v$  and  $vv'w$ .
- A *global  $R$ -overlap* is a local  $R$ -overlap  $uu'vv'w$  such that also  $u R w$ .

- Let  $R$  be the chain relation  $\overline{R}$  or the cyclic relation  $\mathring{R}$ . An  $R$ -repetition of order  $k$  is globally (resp. locally)  *$n$ -avoidable* if there exists an infinite word  $w$  over the alphabet  $\mathcal{A}_n$  such that *each letter* of the alphabet  $\mathcal{A}_n$  occurs *infinitely many times* in  $w$  and  $w$  is globally (resp. locally)  $(R_n, k)$ -free.



- Let  $R$  be the chain relation  $\overline{R}$  or the cyclic relation  $\mathring{R}$ . An  $R$ -repetition of order  $k$  is globally (resp. locally)  *$n$ -avoidable* if there exists an infinite word  $w$  over the alphabet  $\mathcal{A}_n$  such that *each letter* of the alphabet  $\mathcal{A}_n$  occurs *infinitely many times* in  $w$  and  $w$  is globally (resp. locally)  $(R_n, k)$ -free.
- An  $R_{\uparrow}$ -repetition of order  $k$  is globally (resp. locally)  *$n$ -avoidable* if there exists an infinite word  $w$  over the alphabet  $\mathcal{A}_n \cup \{\diamond\}$  such that *each letter* of the alphabet  $\mathcal{A}_n \cup \{\diamond\}$  occurs *infinitely many times* in  $w$  and  $w$  is globally (resp. locally)  $(R_{\uparrow}, k)$ -free.

### Definition

Let  $R$  be the chain relation  $\overline{R}$ , the cyclic relation  $\mathring{R}$  or the relation  $R_{\uparrow}$ .

- The *global avoidability index*  $\gamma(R, k)$  is the minimal  $n$  (if it exists) such that  $R$ -repetitions of order  $k$  are globally  $n$ -avoidable.

### Definition

Let  $R$  be the chain relation  $\overline{R}$ , the cyclic relation  $\mathring{R}$  or the relation  $R_{\uparrow}$ .

- The *global avoidability index*  $\gamma(R, k)$  is the minimal  $n$  (if it exists) such that  $R$ -repetitions of order  $k$  are globally  $n$ -avoidable.
- The *local avoidability index*  $\lambda(R, k)$  is the minimal  $n$  (if it exists) such that  $R$ -repetitions of order  $k$  are locally  $n$ -avoidable.

### Definition

Let  $R$  be the chain relation  $\overline{R}$ , the cyclic relation  $\mathring{R}$  or the relation  $R_{\uparrow}$ .

- The *global avoidability index*  $\gamma(R, k)$  is the minimal  $n$  (if it exists) such that  $R$ -repetitions of order  $k$  are globally  $n$ -avoidable.
- The *local avoidability index*  $\lambda(R, k)$  is the minimal  $n$  (if it exists) such that  $R$ -repetitions of order  $k$  are locally  $n$ -avoidable.
- The indices  $\gamma(R, k^+)$  and  $\lambda(R, k^+)$  are defined as above by replacing  $k$  by  $k^+$ .

- $\lambda(R, k) \geq \gamma(R, k)$

# Relational repetition-freeness

Avoidability index

- $\lambda(R, k) \geq \gamma(R, k)$
- $\lambda(R_{\uparrow}, k) \geq \gamma(R_{\uparrow}, k) \geq 2$

# Relational repetition-freeness

## Avoidability index

- $\lambda(R, k) \geq \gamma(R, k)$
- $\lambda(R_{\uparrow}, k) \geq \gamma(R_{\uparrow}, k) \geq 2$
- $\lambda(\overline{R}, k) \geq \gamma(\overline{R}, k) \geq 3$

# Relational repetition-freeness

Avoidability index

- $\lambda(R, k) \geq \gamma(R, k)$
- $\lambda(R_{\uparrow}, k) \geq \gamma(R_{\uparrow}, k) \geq 2$
- $\lambda(\overline{R}, k) \geq \gamma(\overline{R}, k) \geq 3$
- $\lambda(\mathring{R}, k) \geq \gamma(\mathring{R}, k) \geq 4$



- $\lambda(R, k) \geq \gamma(R, k)$
- $\lambda(R_{\uparrow}, k) \geq \gamma(R_{\uparrow}, k) \geq 2$
- $\lambda(\overline{R}, k) \geq \gamma(\overline{R}, k) \geq 3$
- $\lambda(\mathring{R}, k) \geq \gamma(\mathring{R}, k) \geq 4$
- If  $R$ -repetitions of order  $k$  are  $n$ -avoidable for  $R = \overline{R}$ ,  $R = \mathring{R}$  or  $R = R_{\uparrow}$ , then they are also  $(n + 1)$ -avoidable.

# Avoidability results

	$\overline{R}$		$\mathring{R}$		$R_{\uparrow}$	
	$\gamma$	$\lambda$	$\gamma$	$\lambda$	$\gamma$	$\lambda$
2	6	6	7	7	-	-
$2^+$	4	4	5	5	3	-
3	3	4	4	5	2	-
$3^+$	3	3	4	4	2	2

# Square-freeness and overlap-freeness

	$\overline{R}$		$\mathring{R}$		$R_{\uparrow}$	
	$\gamma$	$\lambda$	$\gamma$	$\lambda$	$\gamma$	$\lambda$
2	6	6	7	7	-	-
$2^+$	4	4	5	5	3	-
3	3	4	4	5	2	-
$3^+$	3	3	4	4	2	2

- One cannot avoid *trivial squares*  $a\Diamond$  or  $\Diamond a$ , where  $a$  is a letter.
- One cannot avoid *local cubes*  $a\Diamond b$ , where  $a$  and  $b$  are letters.

# Square-freeness and overlap-freeness

	$\overline{R}$		$\dot{R}$		$R_{\uparrow}$	
	$\gamma$	$\lambda$	$\gamma$	$\lambda$	$\gamma$	$\lambda$
2	6	6	7	7	-	-
$2^+$	4	4	5	5	3	-
3	3	4	4	5	2	-
$3^+$	3	3	4	4	2	2

- There exists uncountably many infinite words with an infinite number of holes over a three-letter alphabet such that they do not contain any squares other than the trivial ones (Halava, Harju, Kärki'08; Blanchet-Sadri, Mercaş, Scott'09).

# Square-freeness and overlap-freeness

	$\overline{R}$		$\mathring{R}$		$R_{\uparrow}$	
	$\gamma$	$\lambda$	$\gamma$	$\lambda$	$\gamma$	$\lambda$
2	6	6	7	7	-	-
$2^+$	4	4	5	5	3	-
3	3	4	4	5	2	-
$3^+$	3	3	4	4	2	2

- There exists uncountably many infinite words with an infinite number of holes over a three-letter alphabet such that they do not contain any squares other than the trivial ones (Halava, Harju, Kärki'08; Blanchet-Sadri, Mercaş, Scott'09).
- $\gamma(R_{\uparrow}, 2^+) > 2$  (BMS'09; HHK, Séébold'09)

# Square-freeness and overlap-freeness

	$\overline{R}$		$\mathring{R}$		$R_{\uparrow}$	
	$\gamma$	$\lambda$	$\gamma$	$\lambda$	$\gamma$	$\lambda$
2	6	6	7	7	-	-
$2^+$	4	4	5	5	3	-
3	3	4	4	5	2	-
$3^+$	3	3	4	4	2	2

- There exists uncountably many infinite words with an infinite number of holes over a three-letter alphabet such that they do not contain any squares other than the trivial ones (Halava, Harju, Kärki'08; Blanchet-Sadri, Mercaş, Scott'09).
- $\gamma(R_{\uparrow}, 2^+) > 2$  (BMS'09; HHK, Séébold'09)

Theorem (Blanchet-Sadri, Mercaş, Scott'09)

$$\gamma(R_{\uparrow}, 2^+) = 3$$

# Square-freeness and overlap-freeness

	$\overline{R}$		$\dot{R}$		$R_{\uparrow}$	
	$\gamma$	$\lambda$	$\gamma$	$\lambda$	$\gamma$	$\lambda$
2	6	6	7	7	-	-
$2^+$	4	4	5	5	3	-
3	3	4	4	5	2	-
$3^+$	3	3	4	4	2	2

## Theorem (K.'12)

$$\begin{aligned} \gamma(\overline{R}, 2) &= \lambda(\overline{R}, 2) = 6, & \gamma(\dot{R}, 2) &= \lambda(\dot{R}, 2) = 7, \\ \gamma(\overline{R}, 2^+) &= \lambda(\overline{R}, 2^+) = 4, & \gamma(\dot{R}, 2^+) &= \lambda(\dot{R}, 2^+) = 5. \end{aligned}$$

# Cube-freeness and $3^+$ -freeness

	$\overline{R}$		$\mathring{R}$		$R_{\uparrow}$	
	$\gamma$	$\lambda$	$\gamma$	$\lambda$	$\gamma$	$\lambda$
2	6	6	7	7	-	-
$2^+$	4	4	5	5	3	-
3	3	4	4	5	2	-
$3^+$	3	3	4	4	2	2

## Theorem (Manea, Mercaş'07)

*There exist infinitely many cube-free infinite partial words over a binary alphabet containing an infinite number of holes.*

## Corollary

*For any  $k \geq 3$ , we have*

$$\gamma(R_{\uparrow}, k) = 2, \quad \gamma(\overline{R}, k) = 3 \quad \text{and} \quad \gamma(\mathring{R}, k) = 4 .$$



# Cube-freeness and $3^+$ -freeness

	$\overline{R}$		$\mathring{R}$		$R_{\uparrow}$	
	$\gamma$	$\lambda$	$\gamma$	$\lambda$	$\gamma$	$\lambda$
2	6	6	7	7	-	-
$2^+$	4	4	5	5	3	-
3	3	4	4	5	2	-
$3^+$	3	3	4	4	2	2

- For any letters  $x$  and  $y$ , the factor  $x1y$  is a local  $\overline{R}_3$ -cube

# Cube-freeness and $3^+$ -freeness

	$\overline{R}$		$\mathring{R}$		$R_{\uparrow}$	
	$\gamma$	$\lambda$	$\gamma$	$\lambda$	$\gamma$	$\lambda$
2	6	6	7	7	-	-
$2^+$	4	4	5	5	3	-
3	3	4	4	5	2	-
$3^+$	3	3	4	4	2	2

- For any letters  $x$  and  $y$ , the factor  $x1y$  is a local  $\overline{R}_3$ -cube
- Local  $\overline{R}$ -overlaps are 4-avoidable

# Cube-freeness and $3^+$ -freeness

	$\overline{R}$		$\mathring{R}$		$R_{\uparrow}$	
	$\gamma$	$\lambda$	$\gamma$	$\lambda$	$\gamma$	$\lambda$
2	6	6	7	7	-	-
$2^+$	4	4	5	5	3	-
3	3	4	4	5	2	-
$3^+$	3	3	4	4	2	2

- For any letters  $x$  and  $y$ , the factor  $x1y$  is a local  $\overline{R}_3$ -cube
- Local  $\overline{R}$ -overlaps are 4-avoidable

$$\Rightarrow \lambda(\overline{R}, 3) = 4$$

# Cube-freeness and $3^+$ -freeness

	$\overline{R}$		$\mathring{R}$		$R_{\uparrow}$	
	$\gamma$	$\lambda$	$\gamma$	$\lambda$	$\gamma$	$\lambda$
2	6	6	7	7	-	-
$2^+$	4	4	5	5	3	-
3	3	4	4	5	2	-
$3^+$	3	3	4	4	2	2

- Assume that there exists a locally  $(\mathring{R}_4, 3)$ -free word  $w$  containing infinitely many occurrences of each letter in  $\mathcal{A}_4$ .

# Cube-freeness and $3^+$ -freeness

	$\overline{R}$		$\mathring{R}$		$R_{\uparrow}$	
	$\gamma$	$\lambda$	$\gamma$	$\lambda$	$\gamma$	$\lambda$
2	6	6	7	7	-	-
$2^+$	4	4	5	5	3	-
3	3	4	4	5	2	-
$3^+$	3	3	4	4	2	2

- Assume that there exists a locally  $(\mathring{R}_4, 3)$ -free word  $w$  containing infinitely many occurrences of each letter in  $\mathcal{A}_4$ .
- Let  $\mathcal{A} = \{0, 2\}$  or  $\mathcal{B} = \{1, 3\}$ .

# Cube-freeness and $3^+$ -freeness

	$\overline{R}$		$\mathring{R}$		$R_{\uparrow}$	
	$\gamma$	$\lambda$	$\gamma$	$\lambda$	$\gamma$	$\lambda$
2	6	6	7	7	-	-
$2^+$	4	4	5	5	3	-
3	3	4	4	5	2	-
$3^+$	3	3	4	4	2	2

- Assume that there exists a locally  $(\mathring{R}_4, 3)$ -free word  $w$  containing infinitely many occurrences of each letter in  $\mathcal{A}_4$ .
- Let  $\mathcal{A} = \{0, 2\}$  or  $\mathcal{B} = \{1, 3\}$ .
- WLOG: There is a factor 201331 or 201311 in  $w$ .  
It is followed only by letters in  $\mathcal{B}$ : a contradiction.

# Cube-freeness and $3^+$ -freeness

	$\overline{R}$		$\mathring{R}$		$R_{\uparrow}$	
	$\gamma$	$\lambda$	$\gamma$	$\lambda$	$\gamma$	$\lambda$
2	6	6	7	7	-	-
$2^+$	4	4	5	5	3	-
3	3	4	4	5	2	-
$3^+$	3	3	4	4	2	2

- Assume that there exists a locally  $(\mathring{R}_4, 3)$ -free word  $w$  containing infinitely many occurrences of each letter in  $\mathcal{A}_4$ .
- Let  $\mathcal{A} = \{0, 2\}$  or  $\mathcal{B} = \{1, 3\}$ .
- WLOG: There is a factor 201331 or 201311 in  $w$ .  
It is followed only by letters in  $\mathcal{B}$ : a contradiction.
- Local  $\mathring{R}$ -overlaps are 5-avoidable

# Cube-freeness and $3^+$ -freeness

	$\overline{R}$		$\mathring{R}$		$R_{\uparrow}$	
	$\gamma$	$\lambda$	$\gamma$	$\lambda$	$\gamma$	$\lambda$
2	6	6	7	7	-	-
$2^+$	4	4	5	5	3	-
3	3	4	4	5	2	-
$3^+$	3	3	4	4	2	2

- Assume that there exists a locally  $(\mathring{R}_4, 3)$ -free word  $w$  containing infinitely many occurrences of each letter in  $\mathcal{A}_4$ .
- Let  $\mathcal{A} = \{0, 2\}$  or  $\mathcal{B} = \{1, 3\}$ .
- WLOG: There is a factor 201331 or 201311 in  $w$ .  
It is followed only by letters in  $\mathcal{B}$ : a contradiction.
- Local  $\mathring{R}$ -overlaps are 5-avoidable  
 $\Rightarrow \lambda(\mathring{R}, 3) = 5$



# Cube-freeness and $3^+$ -freeness

	$\overline{R}$		$\mathring{R}$		$R_{\uparrow}$	
	$\gamma$	$\lambda$	$\gamma$	$\lambda$	$\gamma$	$\lambda$
2	6	6	7	7	-	-
$2^+$	4	4	5	5	3	-
3	3	4	4	5	2	-
$3^+$	3	3	4	4	2	2

## Theorem

For  $k > 3$ , we have

$$\lambda(R_{\uparrow}, k) = 2, \quad \lambda(\overline{R}, k) = 3 \quad \text{and} \quad \lambda(\mathring{R}, k) = 4.$$

The proof is based on a modification of the Thue-Morse word  $\tau^{\omega}(a)$ , where  $\tau^5(a)$  is replaced by the word  $abbabaabbaaba \diamond babaababbaabbabaab$ .

# Avoidability results

	$\overline{R}$		$\mathring{R}$		$R_{\uparrow}$	
	$\gamma$	$\lambda$	$\gamma$	$\lambda$	$\gamma$	$\lambda$
2	6	6	7	7	-	-
$2^+$	4	4	5	5	3	-
3	3	4	4	5	2	-
$3^+$	3	3	4	4	2	2

Thank you for your attention!